

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 July 2004 (29.07.2004)

PCT

(10) International Publication Number
WO 2004/062624 A3

- (51) International Patent Classification⁷: A61K 31/551, 31/519, 31/535, 31/135
- (21) International Application Number:
PCT/US2004/000809
- (22) International Filing Date: 13 January 2004 (13.01.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/440,076 13 January 2003 (13.01.2003) US
60/492,478 4 August 2003 (04.08.2003) US
- (71) Applicant (for all designated States except US): DYN-
GEN PHARMACEUTICALS, INC. [US/US]; 52 Second
Avenue, Waltham, MA 02451 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): LANDAU, Steven,
B. [US/US]; 44 Tanglewood Road, Wellesley, MA 02481
(US). MILLER, Cheryl, L. [US/US]; 39 Florence Street,
Natick, MA 01760 (US). THOR, Karl, Bruce [US/US];
109 Draymore Way, Morrisville, NC 27560 (US).
- (74) Agents: WENDLER, Helen, E. et al.; Hamilton, Brook,
Smith & Reynolds, P.C.; 530 Virginia Road, P.O. Box 9133,
Concord, MA 01742-9133 (US).
- (81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.
- (84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- (88) Date of publication of the international search report:
7 April 2005
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: METHOD OF TREATING NAUSEA, VOMITING, RETCHING OR ANY COMBINATION THEREOF

(57) Abstract: The invention relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need of treatment. The method comprises administering to a subject in need of treatment a therapeutically effective amount of a compound that has 5-HT₃ receptor antagonist activity and NorAdrenaline Reuptake Inhibitor (NARI) activity. The invention further relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need thereof, comprising coadministering to said subject a first amount of a 5-HT₃ antagonist and a second amount of a NARI, wherein the first and second amounts together comprise a therapeutically effective amount or are each present in a therapeutically effective amount. In addition, the method of the invention comprises administering a NARI alone.

WO 2004/062624 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/00809

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/551, 31/519, 31/535, 31/135

US CL : 514/218, 252.16, 231.2, 649, 653

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/218, 252.16, 231.2, 649, 653

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4,695,568 (NINOMIYA et al) 22 September 1987 (22.09.1987), see entire documents.	1-21
Y	WO 00/061160 A1 (SANOFI-SYNTHELABO) 10 February 2000 (10.02.2000), see entire documents, especially page 6, line 5. and claim 4.	1-21
Y	JP 06016557 A2 (MITSUBISHI CHEMICAL INDUSTRIES) 25 January 1994 (25.01.1994), see entire documents.	1-21
Y,P	US 2004/0048874 A1 (BARDSLEY et al) 11 March 2004 (11.03.2004), see entire documents.	1-21

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

01 November 2004 (01.11.2004)

Date of mailing of the international search report

13 JAN 2005

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (703) 305-3230

Authorized officer

Brian Kwon

Telephone No. 703-308-1235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/00809

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claim Nos.: 22-70
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Please See Continuation Sheet
3. ☐ Claim Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

PCT/US04/00809

Continuation of Box I Reason 2:

Claims 22-70 relate to an extremely large number of possible combinations comprising "5-HT₃ receptor antagonist" and "a noradrenaline reuptake inhibitor". Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compositions claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely "5-HT₃ receptor antagonist" represented by Formula I and examples in page 21, and a "noradrenaline reuptake inhibitor" exemplified in page 28.

Continuation of B. FIELDS SEARCHED Item 3:

STN Online

search terms: nausea, emesis, vomiting, radiation, cancer chemotherapy, vertigo, headache

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 July 2004 (29.07.2004)

PCT

(10) International Publication Number
WO 2004/062624 A2

- (51) International Patent Classification⁷: **A61K**
- (21) International Application Number:
PCT/US2004/000809
- (22) International Filing Date: 13 January 2004 (13.01.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/440,076 13 January 2003 (13.01.2003) US
60/492,478 4 August 2003 (04.08.2003) US
- (71) Applicant (for all designated States except US): **DYNOGEN PHARMACEUTICALS, INC.** [US/US]; 31 St James Avenue, Suite 905, Boston, MA 02116 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **LANDAU, Steven, B.** [US/US]; 44 Tanglewood Road, Wellesley, MA 02481 (US). **MILLER, Cheryl, L.** [US/US]; 39 Florence Street, Natick, MA 01760 (US). **THOR, Karl, Bruce** [US/US]; 109 Draymore Way, Morrisville, NC 27560 (US).
- (74) Agents: **WENDLER, Helen, E. et al.**; Hamilton, Brook, Smith & Reynolds, P.C., 530 Virginia Road, P.O. Box 9133, Concord, MA 01742-9133 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2004/062624 A2

(54) Title: METHOD OF TREATING NAUSEA, VOMITING, RETCHING OR ANY COMBINATION THEREOF

(57) Abstract: The invention relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need of treatment. The method comprises administering to a subject in need of treatment a therapeutically effective amount of a compound that has 5-HT₃ receptor antagonist activity and NorAdrenaline Reuptake Inhibitor (NARI) activity. The invention further relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need thereof, comprising coadministering to said subject a first amount of a 5-HT₃ antagonist and a second amount of a NARI, wherein the first and second amounts together comprise a therapeutically effective amount or are each present in a therapeutically effective amount. In addition, the method of the invention comprises administering a NARI alone.

-1-

METHOD OF TREATING NAUSEA, VOMITING, RETCHING OR ANY COMBINATION THEREOF

RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No.

- 5 60/492,478 filed on August 4, 2003 and U.S. Provisional Application No.
60/440,076 filed on January 13, 2003. The entire teachings of the above
applications are incorporated herein by reference.

BACKGROUND OF THE INVENTION

- Emesis is the act of vomiting and can be described as the forceful expulsion
10 of gastrointestinal contents through the mouth brought about by the descent of the
diaphragm and powerful contractions of the abdominal muscles. Emesis is usually,
but not always, preceded by nausea (the unpleasant feeling that one is about to
vomit). Retching or dry heaves involves the same physiological mechanisms as
vomiting, but occurs against a closed glottis, which prohibits the expulsion of gastric
15 contents. Vomiting, nausea, retching or combinations thereof can be caused by a
number of factors including, but not limited to, anesthetics, radiation, cancer
chemotherapeutic agents, toxic agents, odors, medicines, for example, serotonin
reuptake inhibitors, analgesics such as morphine, antibiotics and antiparasitic agents,
pregnancy and motion. Conditions which are associated with vertigo (e.g.,
20 Meniere's disease and vestibular neuronitis) can also cause nausea, vomiting,
retching or any combination thereof. Headache, caused by, for example, migraine,
increased intracranial pressure or cerebral vascular hemorrhage can also result in
nausea, vomiting, retching or any combination thereof. In addition, certain maladies
of the gastrointestinal (GI) tract, for example, cholecystitis, choledocholithiasis,
25 intestinal obstruction, acute gastroenteritis, perforated viscus, dyspepsia resulting
from, for example, gastroesophageal reflux disease, peptic ulcer disease,
gastroparesis, gastric or esophageal neoplasms, infiltrative gastric disorders (e.g.,
Menetrier's syndrome, Crohn's disease, eosinophilic gastroenteritis, sarcoidosis and
amyloidosis), gastric infections (e.g., CMV, fungal, TB and syphilis), parasites (e.g.,

-2-

Giardia lamblia and *Strongyloides stercoralis*), chronic gastric volvulus, chronic intestinal ischemia, altered gastric motility disorders and/or food intolerance or Zollinger-Ellison syndrome can result in vomiting, nausea, retching or any combination thereof. However, in some cases of vomiting, nausea, retching or any combination thereof, no etiology can be determined despite extensive diagnostic testing (e.g., Cyclic Vomiting Syndrome).

Nausea, vomiting and retching are defined as acute when symptoms are present for less than a week. The causes of nausea, vomiting and retching of short duration are often separable from etiologies leading to more chronic symptoms.

10 Nausea, vomiting and retching are defined as chronic when symptoms are present for over a week. For example, symptoms can be continuous or intermittent and last for months or years.

The vomiting reflex is triggered by stimulation of chemoreceptors in the upper GI tract and mechanoreceptors in the wall of the GI tract which are activated by both contraction and distension of the gut as well as by physical damage. A coordinating center in the central nervous system controls the emetic response. This center is located in the paraventricular reticular formation in the lateral medullary region of the brain. Afferent nerves to the vomiting center arise from abdominal splanchnic and vagal nerves, vestibulo-labyrinthine receptors, the cerebral cortex and the chemoreceptor trigger zone (CTZ). The CTZ lies adjacent in the area postrema and contains chemoreceptors that sample both blood and cerebrospinal fluid. Direct links exist between the emetic center and the CTZ. The CTZ is exposed to emetic stimuli of endogenous origin (e.g., hormones) as well as to stimuli of exogenous origin, such as drugs. The efferent branches of cranial nerves V, VII and IX, as well as the vagus nerve and sympathetic trunk produce the complex coordinated set of muscular contractions, cardiovascular responses and reverse peristalsis that characterize vomiting.

Of significant clinical relevance is the nausea and vomiting resulting from the administration of general anesthetics (commonly referred to as, post-operative nausea and vomiting, PONV), chemotherapeutic agents and radiation therapy.

In fact, the symptoms caused by the chemotherapeutic agents can be so severe that the patient refuses further treatment. Three types of emesis are associated with the use of chemotherapeutic agents. The first, is acute emesis, which occurs within the first 24 hours of chemotherapy. The second, is delayed emesis
5 which occurs 24 hours or more after chemotherapy administration. The third, is anticipatory emesis, which begins prior to the administration of chemotherapy, usually in patients whose emesis was poorly controlled during a previous chemotherapy cycle.

PONV is also an important patient problem and one that patients rate as the
10 most distressing aspect of operative procedure, even above pain. Consequently, the need for an effective anti-emetic in this area is important. As a clinical problem PONV is troublesome and requires staff around to ensure that vomitus is not regurgitated, which can have very serious clinical sequelae. Further, there are certain operative procedures where it is clinically important that patients do not
15 vomit. For example, in ocular surgery where intra-cranial ocular pressure can increase to the extent that stitches are ruptured and the operative procedure is set back in terms of success to a marked degree.

There are a number groups of agents that are used clinically for the treatment of emesis. These groups include: anticholinergics, antihistamines, phenothiazines,
20 butyrophenones, cannabinoids, benzamides, glucocorticoids, benzodiazepines and 5-HT₃ receptor antagonists. In addition, tricyclic antidepressants have also been used on a limited basis.

The phenothiazines, which include prochlorperazine and chlorpromazine, block dopamine type-2 receptors in the CTZ. However, the side effects, for
25 example, extrapyramidal symptoms, such as, dystonia and akathisia, sedation, anticholinergic effect and orthostatic hypotension make the use of the phenothiazines a less than desirable therapy.

Anticholinergics used in the treatment of nausea and vomiting, include scopolamine (e.g., in treating motion sickness). However, drowsiness is a
30 significant side effect.

Antihistamines (dimenhydrinate and diphenhydramine) are mainly used for motion sickness and in antiemetic combinations to reduce extrapyramidal side effects of dopamine receptor antagonists. As a single agent, the antihistamines have modest antiemetic activity and include sedation and anticholinergic effects as the major drawbacks.

Butyrophenones, for example, haloperidol and droperidol, work by blocking dopamine receptors in the CTZ. The side effects of butyrophenones include akathisia, dystonia and hypotension.

Cannabinoids such as tetrahydrocannabinol and nabilone have shown limited efficacy (see, e.g. Sallan *et al.*, *N. Eng. J. Med.*, 302: 135-138 (1980)). In addition, the side effects include euphoria, dizziness, paranoid ideation and somnolence.

Benzamides include, for example, metoclopramide, cisapride and trimethobenzamide. However, side effects which include extrapyramidal symptoms and diarrhea make the use of benzamides a less than desirable therapy.

Benzodiazapines include, for example, lorazepam. Side effects of the benzodiazapines include perceptual disturbances, urinary incontinence, hypotension, diarrhea, sedation and amnesia.

Corticosteroids such as dexamethasone and methylprednisolone are useful in combination therapy, but shown little efficacy as a single agent. Side effects include, hyperglycemia, euphoria, insomnia and rectal pain.

The antiemetic property of tricyclic antidepressants has been assessed on a limited basis (see, e.g., Prakash *et al.*, *Dig. Dis. Sci.* 43(9):1951-1956 (1998)) and cyclic vomiting syndrome (Prakash and Clouse, *Am. J. Gastroenterol.*, 94(10): 2855-2860 (1999)).

However, the undesirable side effects associated with the use of tricyclic antidepressants are a significant drawback for this therapy. For example, the anticholinergic properties of the tricyclic antidepressants can cause dry mouth, constipation, blurred vision, urinary retention, weight gain, hypertension and cardiac side effects, such as palpitations and arrhythmia.

Antagonism of the 5-HT₃ receptor has been the focus of antiemetic therapy. More specifically, 5-HT₃ receptors are widely distributed in the mammalian central,

peripheral and enteric nervous systems. The enteric nervous system resides within the walls of the gastrointestinal tract. 5-HT₃ receptors have been found to play an important role in the control of vomiting in a variety of mammals including humans (Veyrat-Follet *et al.*, *Drugs* 53(2):206-234 (1997)). The receptors are present in the

5 part of the brain that is involved in controlling vomiting as well as in the gastrointestinal tract. Receptors at both locations have been shown to be involved in vomiting. It is thought that 5-HT released from the enterochromaffin cells of the gastrointestinal mucosa acts on 5-HT₃ receptors to initiate the vomiting reflex. Chemotherapy and radiotherapy, two important clinical causes of vomiting, can

10 cause release of 5-HT from the enterochromaffin cells. Chemotherapeutic agents also appear to act directly on the chemoreceptor trigger zone (CTZ) of the vomiting center in the brain that then feeds onto neurons containing 5-HT₃ receptors to initiate vomiting. That is, activation of the chemoreceptor trigger zone (CTZ) triggers the release of neurotransmitters that activate the vomiting center. CTZ neurotransmitters

15 that are thought to cause emesis include, but are not limited to, dopamine, serotonin, histamine and norepinephrine.

However, improved treatment regimens are still needed. For example, the use of 5-HT₃ receptor antagonists such as ondansetron, granisetron and tropisetron has been shown to be less effective for delayed nausea and vomiting than for acute

20 symptoms. In addition, efficacy of the 5-HT₃ receptor antagonists appears to be less pronounced for moderate emetogenic chemotherapy regimens than for cisplatin-containing regimens. Further, control over nausea appears to be significantly less than control over vomiting. Further, the efficacy of the agents appears to diminish across repeated days and across repeated chemotherapy cycles (see, e.g., Morrow *et al.*, *Cancer* 76(3): 343-357 (1995)).

25

As such, improved methods for the treatment of vomiting, nausea, retching or any combination thereof are needed.

SUMMARY OF THE INVENTION

The invention relates to a method of treating nausea, vomiting, retching or

30 any combination thereof in a subject in need of treatment. The method comprises


-6-

administering to a subject in need of treatment a therapeutically effective amount of a compound that has 5-HT₃ receptor antagonist activity and NorAdrenaline Reuptake Inhibitor (NARI) activity.

In a particular embodiment, the compounds having 5-HT₃ receptor antagonist activity and NARI activity are thieno[2,3-d]pyrimidine derivatives such as those described in U.S. Patent No. 4,695,568, the entire content of which is incorporated herein by reference.

In a specific embodiment, the compounds having 5-HT₃ receptor antagonist activity and NARI activity are represented by structural Formula I:

10




wherein, R₁ and R₂ independently represent hydrogen, halogen or a C₁-C₆ alkyl group; or R₁ and R₂ together with the carbon atoms to which they are attached form a cycloalkylene group having 5 to 6 carbon atoms;

15

R₃ and R₄ independently represent hydrogen or a C₁-C₆ alkyl group;

R₅ is hydrogen, C₁-C₆ alkyl,



or -C(O)-NH-R₆,


20

wherein m is an integer from about 1 to about 3, X is halogen and R₆ is a C₁-C₆ alkyl group;

Ar is a substituted or unsubstituted phenyl, 2-thienyl or 3-thienyl group; and n is 2 or 3; or a pharmaceutically acceptable salt thereof.

-7-

In a specific embodiment, the compound having 5-HT₃ receptor antagonist activity and NARI activity is represented by the formula:



5 or a pharmaceutically acceptable salt thereof. This compound is referred to as MCI-225 of DDP-225. The chemical name of the structure set forth in the formula is: 4-(2-fluorophenyl)-6-methyl-2-(1-piperazinyl)thieno[2,3-d]pyrimidine.

10 In one embodiment, the nausea, vomiting, retching or any combination thereof can be caused by anesthetics, radiation, cancer chemotherapeutic agents, toxic agents, odors, medicines, pregnancy and motion.

In a particular embodiment, the nausea, vomiting, retching or any combination thereof can be caused by administration of general anesthetics associated with surgical procedures.

15 In a more particular embodiment, the nausea, vomiting, retching or any combination thereof can be caused by administration of chemotherapeutic agents, radiation therapy or a combination thereof.

In yet another embodiment, the nausea, vomiting, retching or any combination thereof can be caused by conditions which are associated with vertigo. For example, the nausea, vomiting, retching or any combination thereof can be
20 caused by Meniere's disease or vestibular neuronitis.

In another embodiment, the nausea, vomiting, retching or any combination thereof can be caused by headache. In particular embodiments, the headache is a result of migraine, increased intracranial pressure or cerebral vascular hemorrhage.

In yet another embodiment, the nausea, vomiting, retching or any combination thereof can be caused by maladies of the gastrointestinal (GI) tract. In a particular embodiment, the malady of the gastrointestinal tract is selected from the group consisting of cholecystitis, choledocholithiasis, intestinal obstruction, acute gastroenteritis, perforated viscus, dyspepsia and Zollinger-Ellison syndrome.

In a further embodiment, the vomiting, nausea, retching or any combination thereof, can be of undetermined etiology. In a particular embodiment, the nausea, vomiting, retching or any combination thereof can be characterized as Cyclic Vomiting Syndrome.

10 The invention further relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need thereof, comprising coadministering to said subject a therapeutically effective amount of a 5-HT₃ receptor antagonist and a therapeutically effective amount of a NARI.

The invention further relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need thereof, comprising coadministering to said subject a first amount of a 5-HT₃ receptor antagonist and a second amount of a NARI, wherein the first and second amounts together comprise a therapeutically effective amount.

20 In one embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof caused by anesthetics, radiation, cancer chemotherapeutic agents, toxic agents, odors, medicines, pregnancy and motion.

In a particular embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof caused by administration of general anesthetics associated with surgical procedures.

25 In a more particular embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof caused by administration of chemotherapeutic agents, radiation therapy or a combination thereof.

In yet another embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof caused by conditions which are

-9-

associated with vertigo. For example, the nausea, vomiting, retching or any combination thereof can be caused by Meniere's disease or vestibular neuronitis.

In another embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof caused by headache. In particular
5 embodiments, the headache is a result of migraine, increased intracranial pressure or cerebral vascular hemorrhage.

In yet another embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof caused by maladies of the gastrointestinal (GI) tract. In a particular embodiment, the malady of the
10 gastrointestinal tract is selected from the group consisting of cholecystitis, choledocholithiasis, intestinal obstruction, acute gastroenteritis, perforated viscus, dyspepsia and Zollinger-Ellison syndrome.

In a further embodiment, the coadministration can be used to treat vomiting, nausea, retching or any combination thereof, of undetermined etiology. In a
15 particular embodiment, the nausea, vomiting, retching or any combination thereof can be characterized as Cyclic Vomiting Syndrome.

In addition, the invention relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need thereof comprising administering a therapeutically effective amount of a NARI. In this embodiment, the
20 NARI is characterized by the substantial absence of anticholinergic effects.

In one embodiment, the nausea, vomiting, retching or any combination thereof can be caused by anesthetics, radiation, cancer chemotherapeutic agents, toxic agents, odors, medicines, pregnancy and motion.

In a particular embodiment, the nausea, vomiting, retching or any
25 combination thereof can be caused by administration of general anesthetics associated with surgical procedures.

In a more particular embodiment, the nausea, vomiting, retching or any combination thereof can be caused by administration of chemotherapeutic agents, radiation therapy or a combination thereof.

-10-

In yet another embodiment, the nausea, vomiting, retching or any combination thereof can be caused by conditions which are associated with vertigo. For example, the nausea, vomiting, retching or any combination thereof can be caused by Meniere's disease or vestibular neuronitis.

5 In another embodiment, the nausea, vomiting, retching or any combination thereof can be caused by headache. In particular embodiments, the headache is a result of migraine, increased intracranial pressure or cerebral vascular hemorrhage.

In yet another embodiment, the nausea, vomiting, retching or any combination thereof can be caused by maladies of the gastrointestinal (GI) tract. In a
10 particular embodiment, the malady of the gastrointestinal tract can be selected from the group consisting of cholecystitis, choledocholithiasis, intestinal obstruction, acute gastroenteritis, perforated viscus, dyspepsia and Zollinger-Ellison syndrome.

In a further embodiment, the vomiting, nausea, retching or any combination thereof, can be of undetermined etiology. In a particular embodiment, the nausea,
15 vomiting, retching or any combination thereof can be characterized as Cyclic Vomiting Syndrome.

The invention further relates to pharmaceutical compositions useful for the treatment of a nausea, vomiting, retching or any combination thereof. The pharmaceutical composition comprises a first amount of a 5-HT₃ receptor antagonist
20 compound and a second amount of a NARI compound. The pharmaceutical compositions of the present invention can optionally contain a pharmaceutically acceptable carrier. The 5-HT₃ receptor antagonist and the NARI can each be present in the pharmaceutical composition in a therapeutically effective amount. In another aspect, said first and second amounts can together comprise a therapeutically
25 effective amount.

The pharmaceutical composition can be used to treat vomiting, nausea, retching or combinations thereof caused by a number of factors including, but not limited to, anesthetics, radiation, cancer chemotherapeutic agents, toxic agents, odors, medicines, for example, serotonin reuptake inhibitors, analgesics such as
30 morphine, antibiotics and antiparasitic agents, pregnancy and motion. Conditions which are associated with vertigo (e.g., Meniere's disease and vestibular neuronitis)

-11-

can also cause nausea, vomiting, retching or any combination thereof. Headache, caused by, for example, migraine, increased intracranial pressure or cerebral vascular hemorrhage can also result in nausea, vomiting, retching or any combination thereof. In addition, certain maladies of the gastrointestinal (GI) tract, for example,

5 cholecystitis, choledocholithiasis, intestinal obstruction, acute gastroenteritis, perforated viscus, dyspepsia resulting from, for example, gastroesophageal reflux disease, peptic ulcer disease, gastroparesis, gastric or esophageal neoplasms, infiltrative gastric disorders (e.g., Menetrier's syndrome, Crohn's disease, eosinophilic gastroenteritis, sarcoidosis and amyloidosis) gastric infections (e.g.,

10 CMV, fungal, TB and syphilis), parasites (e.g., *Giardia lamblia* and *Strongyloides stercoralis*), chronic gastric volvulus, chronic intestinal ischemia, altered gastric motility and/or food intolerance or Zollinger-Ellison syndrome can result in vomiting, nausea, retching or any combination thereof.

The invention further relates to use of a compound that has 5-HT₃ receptor

15 antagonist activity and NARI activity for the manufacture of a medicament for treating nausea, vomiting, retching or any combination thereof. In addition, the invention also relates to the use of a pharmaceutical composition comprising a first amount of a 5-HT₃ receptor antagonist compound and a second amount of a NARI compound for the manufacture of a medicament for the treatment of a nausea,

20 vomiting, retching or any combination thereof. The pharmaceutical composition used for the manufacture of a medicament for treating nausea, vomiting, retching or any combination thereof can optionally contain a pharmaceutically acceptable carrier. The 5-HT₃ receptor antagonist and the NARI can each be present in the pharmaceutical composition in a therapeutically effective amount or said first and

25 second amounts can together comprise a therapeutically effective amount. Further, the invention relates to the use of a NARI for the manufacture of a medicament for treating nausea, vomiting, retching or any combination thereof.

The foregoing and other objects, features and advantages of the invention will be apparent from the following more particular description of preferred

30 embodiments of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a bar graph of cisplatin-induced retches per hour versus time (hours) post administration of cisplatin in male ferrets treated with a 5 mg/kg dose of cisplatin and vehicle.

5 FIG. 2 is a bar graph of cisplatin-induced vomits per hour versus time (hours) post administration of cisplatin in male ferrets treated with a 5 mg/kg dose of cisplatin and vehicle.

FIG. 3 is a bar graph of the total number of cisplatin-induced emetic events (retches and vomits combined) in male ferrets treated with cisplatin at a dose of 5
10 mg/kg followed by ondansetron at 5 mg/kg or 10 mg/kg or vehicle alone.

FIG. 4 is a bar graph of the effect of ondansetron on the latency of the first cisplatin-induced emetic event (retch or vomit) in male ferrets treated with cisplatin at a dose of 5 mg/kg followed by ondansetron at 5 mg/kg or 10 mg/kg or vehicle alone.

15 FIG. 5 is a bar graph of the total number of cisplatin-induced emetic events (retches and vomits combined) in male ferrets treated with cisplatin at a dose of 5 mg/kg followed by a single dose of MCI-225 (1 mg/kg, 10 mg/kg or 30 mg/kg), two 30 mg/kg doses at 3 hour intervals or vehicle alone.

FIG. 6 is a bar graph of the effect of ondansetron on the latency of the first
20 cisplatin-induced emetic event (retch or vomit) in male ferrets treated with cisplatin at a dose of 5 mg/kg followed by a single dose of MCI-225 (1 mg/kg, 10 mg/kg or 30 mg/kg), two 30 mg/kg doses at 3 hour intervals or vehicle alone.

DETAILED DESCRIPTION OF THE INVENTION

The invention relates to methods of treating nausea, vomiting, retching or
25 any combination thereof.

MONOAMINE NEUROTRANSMITTERS:

Monoamine neurotransmitters such as noradrenaline (also referred to as norepinephrine), serotonin (5-hydroxytryptamine, 5-HT) and dopamine are known and disturbances in these neurotransmitters have been indicated in many types of

-13-

disorders, such as depression. These neurotransmitters travel from the terminal of a neuron across a small gap referred to as the synaptic cleft and bind to receptor molecules on the surface of a second neuron. This binding elicits intracellular changes that initiate or activate a response or change in the postsynaptic neuron.

- 5 Inactivation occurs primarily by transport of the neurotransmitter back into the presynaptic neuron, which is referred to as reuptake. These neurons can be found both in the Central Nervous System (CNS) and in the Enteric Nervous System (ENS).

NORADRENALINE AND NORADRENALINE REUPTAKE INHIBITORS:

- 10 As used herein, the term NorAdrenaline Reuptake Inhibitor (NARI) refers to an agent (e.g., a molecule, a compound) which can inhibit noradrenaline transporter function. For example, a NARI can inhibit binding of a ligand of a noradrenaline transporter to said transporter and/or inhibit transport (e.g., uptake or reuptake of noradrenaline). As such, inhibition of the noradrenaline transport function in a
15 subject, can result in an increase in the concentration of physiologically active noradrenaline. It is understood that NorAdrenergic Reuptake Inhibitor and NorEpinephrine Reuptake Inhibitor (NERI) are synonymous with NorAdrenaline Reuptake Inhibitor (NARI).

- As used herein, noradrenaline transporter refers to naturally occurring
20 noradrenaline transporters (e.g., mammalian noradrenaline transporters (e.g., human (*Homo sapiens*) noradrenaline transporters, murine (e.g., rat, mouse) noradrenaline transporters)) and to proteins having an amino acid sequence which is the same as that of a corresponding naturally occurring noradrenaline transporter (e.g., recombinant proteins). The term includes naturally occurring variants, such as
25 polymorphic or allelic variants and splice variants.

- In certain embodiments, the NARI can inhibit the binding of a ligand (e.g., a natural ligand such as noradrenaline, or other ligand such as nisoxetine) to a noradrenaline transporter. In other embodiments, the NARI can bind to a noradrenaline transporter. For example, in a preferred embodiment, the NARI can
30 bind to a noradrenaline transporter, thereby inhibiting binding of a ligand to said

transporter and inhibiting transport of said ligand. In another preferred embodiment, the NARI can bind to a noradrenaline transporter, and thereby inhibit transport.

The NARI activity of a compound can be determined employing suitable assays. More specifically, to determine the inhibition constant (K_i) for
5 noradrenaline reuptake, an assay which monitors inhibition of noradrenaline (NA) uptake can be used. For example, radiolabelled noradrenaline, such as [^3H]NA and the test compound of interest can be incubated under conditions suitable for uptake with brain tissue or a suitable fraction thereof, for example, a synaptosomal fraction from rat brain tissue (harvested and isolated in accordance with generally accepted
10 techniques), and the amount of uptake of [^3H]NA in the tissue or fraction can be determined (e.g., by liquid scintillation spectrometry). IC_{50} values can be calculated by nonlinear regression analysis. The inhibition constants, K_i values, can then be calculated from the IC_{50} values using the Cheng-Prusoff equation:

$$15 \quad K_i = \frac{\text{IC}_{50}}{1 + ([L]/K_d)}$$

wherein $[L]$ = the concentration of free radioligand used in the assay and K_d = the equilibrium dissociation constant of the radioligand. To determine the non-specific uptake, incubations can be performed by following the same assay, but in the
absence of test compound at 4°C (i.e., under conditions not suitable for uptake).

20 In a preferred embodiment, NARI activity is determined using the radioligand uptake assay described above, according to the procedure detailed in Eguchi *et al.*, *Arzneim.-Forschung/Drug Res.*, 47(12): 1337-47 (1997).

Specifically, rats are decapitated and the cortical, hypothalamic, hippocampal and striatal tissues are rapidly dissected. The tissues are homogenized (Potter
25 homogenizer with Teflon pestle) in 10 volumes of ice cold 0.32 mol/L sucrose. The P_2 fraction is obtained by centrifugation at $1000 \times g$ for 10 minutes and $11500 \times g$ for 20 minutes and suspended in Krebs-Ringer phosphate buffer, pH 7.4 (124 mmol/L NaCl, 5 mmol/L KCl, 20 mmol/L Na_2HPO_4 , 1.2 mmol/L KH_2PO_4 , 1.3

-15-

mmol/L MgSO₄, 0.75 mmol/L CaCl₂, 10 mmol/L glucose). The [³H]NA uptake assays are performed on the cortical and hypothalamic synaptosomes.

The assay tubes contain radiolabeled noradrenaline, [³H]NA, in a volume of 0.2 mL, compounds at 5 or more concentrations in a volume of 0.1 mL, and the oxygenated buffer described above in a volume of 0.5 mL. After 5 minutes preincubation at 37 °C, uptake is initiated by the addition of the synaptosomal fraction in volume of 0.2 mL. The final concentration of [³H]NA in the incubation mixtures is 0.25 µmol/L. The reaction is stopped after 5 minutes by filtration through Whatman GF/B glass fiber filter under a vacuum with a cell harvester. The filter is rinsed three times with 4 mL of saline and placed in a scintillation vial containing 10 mL of Atomlight (Du Pont/NEN Research Products). Radioactivity is measured by liquid scintillation spectrometry. For determination of non-specific uptake, incubations are performed at 4 °C without the addition of test compounds. IC₅₀ values are calculated by nonlinear regression analysis. Inhibitor constants, K_i values, are calculated from the IC₅₀ values using the Cheng-Prusoff equation.

NARI compounds suitable for use in the invention have a K_i value for NARI activity of about 500 nmol/L or less, such as about 250 nmol/L or less, for example, about 100 nmol/L or less. It is preferred that the K_i value for NARI activity be about 100 nmol/L or less. It is understood that the exact value of the K_i for a particular compound can vary depending on the assay conditions employed for determination (e.g., radioligand and tissue source). As such, it is preferred that the NARI activity be assessed essentially according to the radioligand binding assay described in Eguchi *et al.*, *Arzneim.-Forschung/Drug Res.*, 47(12): 1337-47 (1997) and discussed in detail above.

In addition, to possessing sufficient NARI activity, it is preferred that the NARI compounds possess one or more characteristics selected from the group consisting of:

- a. the substantial absence of anticholinergic effects;
- b. the selective inhibition of noradrenaline reuptake as compared to inhibition of serotonin reuptake; and

-16-

- c. the selective inhibition of noradrenaline reuptake as compared to inhibition of dopamine reuptake.

Selective inhibition of noradrenaline reuptake as compared to inhibition of serotonin or dopamine reuptake can be determined by comparing the K_i values for the respective reuptake inhibitions. The inhibition constants for serotonin and dopamine reuptake can be determined as described above for noradrenaline, but employing the appropriate radioligand and tissue for the activity being assessed (e.g., [^3H] 5-HT for serotonin, using e.g., hypothalamic or cortical tissue and [^3H]DA for dopamine (DA), using e.g., striatal tissue).

- 10 A preferred method of determining serotonin reuptake inhibition and dopaminergic reuptake inhibition is described in Eguchi *et al.*, *Arzneim.-Forschung/Drug Res.*, 47(12): 1337-47 (1997). Specifically, rats are decapitated and the cortical, hypothalamic, hippocampal and striatal tissues are rapidly dissected. The tissues are homogenized (Potter homogenizer with Teflon pestle) in 10 volumes of ice cold 0.32 mol/L sucrose. The P_2 fraction is obtained by centrifugation at 1000 x g for 10 minutes and 11500 x g for 20 minutes and suspended in Krebs-Ringer phosphate buffer, pH 7.4 (124 mmol/L NaCl, 5 mmol/L KCl, 20 mmol/L Na_2HPO_4 , 1.2 mmol/L KH_2PO_4 , 1.3 mmol/L MgSO_4 , 0.75 mmol/L CaCl_2 , 10 mmol/L glucose). The [^3H]5-HT uptake assays are performed on the cortical, hypothalamic and
15 of ice cold 0.32 mol/L sucrose. The P_2 fraction is obtained by centrifugation at 1000 x g for 10 minutes and 11500 x g for 20 minutes and suspended in Krebs-Ringer phosphate buffer, pH 7.4 (124 mmol/L NaCl, 5 mmol/L KCl, 20 mmol/L Na_2HPO_4 , 1.2 mmol/L KH_2PO_4 , 1.3 mmol/L MgSO_4 , 0.75 mmol/L CaCl_2 , 10 mmol/L glucose). The [^3H]5-HT uptake assays are performed on the cortical, hypothalamic and
20 hippocampal synaptosomes, and the [^3H]DA uptake assays are performed on striatal synaptosomes.

- The assay tubes contain the appropriate radiolabeled ligand (i.e., [^3H]5-HT or [^3H]DA), in a volume of 0.2 mL, compounds at 5 or more concentrations in a volume of 0.1 mL, and the oxygenated buffer described above in a volume of 0.5
25 mL. After 5 minutes preincubation at 37 °C, uptake is initiated by the addition of the synaptosomal fraction in volume of 0.2 mL. The final concentration of [^3H]DA in the striatal incubation mixtures is 0.4 $\mu\text{mol/L}$. The final concentrations of [^3H]5-HT in the cortical, hypothalamic and hippocampal synaptosome incubation mixtures are 0.02 $\mu\text{mol/L}$, 0.04 $\mu\text{mol/L}$ and 0.08 $\mu\text{mol/L}$. The reaction is stopped after 5
30 minutes ([^3H]5-HT) or 3 minutes [^3H]DA by filtration through Whatman GF/B glass

-17-

fiber filter under a vacuum with a cell harvester. The filter is rinsed three times with 4 mL of saline and placed in a scintillation vial containing 10 mL of Atomlight (Du Pont/NEN Research Products). Radioactivity is measured by liquid scintillation spectrometry. For determination of non-specific uptake incubations are performed at 5 4 °C without the addition of test compounds. IC₅₀ values are calculated by nonlinear regression analysis. Inhibition constants, Ki values, are calculated from the IC₅₀ values using the Cheng-Prusoff equation.

Following determination of the Ki values for inhibition of noradrenaline, serotonin and/or dopamine uptake, the ratio of the activities can be determined.

10 Selective inhibition of noradrenaline reuptake as compared to inhibition of serotonin reuptake and/or dopaminergic reuptake, refers to a compound having a Ki value for inhibition of serotonin (re)uptake and/or dopamine (re)uptake which is about 10 times or more than the Ki for inhibition of noradrenaline (re)uptake. That is, the ratio, Ki inhibition of serotonin (re)uptake / Ki inhibition of noradrenaline
15 (re)uptake, is about 10 or more, such as about 15 or more, about 20 or more, for example, about 30, 40 or 50 or more. Likewise, the ratio, Ki inhibition of dopamine (re)uptake / Ki inhibition noradrenaline (re)uptake, is about 10 or more, such as about 15 or more, about 20 or more, for example, about 30, 40 or 50 or more.

It is preferred that the Ki values for comparison are determined according to
20 the method of Eguchi *et al.*, discussed in detail above. It is most preferred, that the Ki values for NARI activity and inhibition of serotonin reuptake activity, which are compared to determine selective inhibition are determined according to the method of Eguchi *et al.* using a synaptosomal preparation from rat hypothalamic tissue. Further, it is most preferred, that the Ki values for NARI activity and inhibition of
25 dopamine reuptake activity, which are compared to determine selective inhibition are determined according to the method of Eguchi *et al.* using a synaptosomal preparation from rat hypothalamic tissue for inhibition of noradrenaline uptake and from rat striatal tissue for inhibition of dopamine uptake.

In another embodiment, the NARI is characterized by the substantial absence
30 of anticholinergic effects. As used herein, substantial absence of anticholinergic effects, refers to a compound which has an IC₅₀ value for binding to muscarinic

receptors of about 1 $\mu\text{mol/L}$ or more. The IC_{50} value for binding to muscarinic receptors can be determined using a suitable assay, such as an assay which determines the ability of a compound to inhibit the binding of suitable radioligand to muscarinic receptors. A preferred assay for determination of the IC_{50} value for
5 binding of a compound to muscarinic receptors is described in Eguchi *et al.*, *Arzneim.-Forschung/Drug Res.*, 47(12): 1337-47 (1997).

Specifically, the binding assays for determination of binding to muscarinic receptors can be performed on tissue isolated from the rat cerebral cortex. The buffer is any suitable buffer, for example, 50 mmol/L Tris-HCl, pH=7.4. The
10 preferred radiolabeled ligand is [^3H]QNB (3-quinuclidinyl benzilate) which is present in a final concentration of 0.2 nmol/L. The test compound is added at various concentrations and the resulting mixtures are incubated for 60 minutes at 37 $^{\circ}\text{C}$. The reaction is terminated by rapid vacuum filtration onto glass fiber filter. Radioactivity trapped on the filter is measured by scintillation spectrometry. Non-
15 specific binding is determined using 100 $\mu\text{mol/L}$ atropine. IC_{50} values can be calculated by nonlinear regression analysis.

In a particular embodiment, the NARI compound can be selected from venlafaxine, duloxetine, bupropion, milnacipran, reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotiline,
20 viloxazine and atomoxetine.

In a preferred embodiment, the NARI compound can be selected from reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotiline, viloxazine and atomoxetine.

SETOTONIN AND 5-HT₃ RECEPTOR ANTAGONISTS:

25 The neurotransmitter serotonin was first discovered in 1948 and has subsequently been the subject of substantial scientific research. Serotonin, also referred to as 5-hydroxytryptamine (5-HT), acts both centrally and peripherally on discrete 5-HT receptors. Currently, fourteen subtypes of serotonin receptors are recognized and delineated into seven families, 5-HT₁ through 5-HT₇. These
30 subtypes share sequence homology and display some similarities in their specificity

for particular ligands. A review of the nomenclature and classification of the 5-HT receptors can be found in *Neuropharm.*, 33: 261-273 (1994) and *Pharm. Rev.*, 46:157-203 (1994).

5 5-HT₃ receptors are ligand-gated ion channels that are extensively distributed on enteric neurons in the human gastrointestinal tract, as well as other peripheral and central locations. Activation of these channels and the resulting neuronal depolarization have been found to affect the regulation of visceral pain, colonic transit and gastrointestinal secretions. Antagonism of the 5-HT₃ receptors has the potential to influence sensory and motor function in the gut.

10 As used herein, 5-HT₃ receptor refers to naturally occurring 5-HT₃ receptors (e.g., mammalian 5-HT₃ receptors (e.g., human (*Homo sapiens*) 5-HT₃ receptors, murine (e.g., rat, mouse) 5-HT₃ receptors)) and to proteins having an amino acid sequence which is the same as that of a corresponding naturally occurring 5-HT₃ receptor (e.g., recombinant proteins). The term includes naturally occurring
15 variants, such as polymorphic or allelic variants and splice variants.

As used herein, the term 5-HT₃ receptor antagonist refers to an agent (e.g., a molecule, a compound) which can inhibit 5-HT₃ receptor function. For example, a 5-HT₃ receptor antagonist can inhibit binding of a ligand of a 5-HT₃ receptor to said receptor and/or inhibit a 5-HT₃ receptor-mediated response (e.g., reduce the ability
20 of 5-HT₃ to evoke the von Bezold-Jarisch reflex).

In certain embodiments, the 5-HT₃ receptor antagonist can inhibit binding of a ligand (e.g., a natural ligand, such as serotonin (5-HT₃), or other ligand such as GR65630) to a 5-HT₃ receptor. In certain embodiments, the 5-HT₃ receptor antagonist can bind to a 5-HT₃ receptor. For example, in a preferred embodiment,
25 the 5-HT₃ receptor antagonist can bind to a 5-HT₃ receptor, thereby inhibiting the binding of a ligand to said receptor and a 5-HT₃ receptor-mediated response to ligand binding. In another preferred embodiment, the 5-HT₃ receptor antagonist can bind to a 5-HT₃ receptor, and thereby inhibit a 5-HT₃ receptor-mediated response.

5-HT₃ receptor antagonists can be identified and activity assessed by any
30 suitable method, for example, by a method which assesses the ability of a compound to inhibit radioligand binding to 5-HT₃ receptor (see, for example, Eguchi *et al.*,

Arzneim.-Forschung/Drug Res., 47(12): 1337-47 (1997) and G. Kilpatrick *et al.*, *Nature*, 330: 746-748 (1987)) and/or by their effect on the 5-HT₃-induced von Bezold-Jarisch (B-J) reflex in the cat or rat (following the general methods described by Butler *et al.*, *Br. J. Pharmacol.*, 94: 397-412 (1988) and Ito *et al.*, *J. Pharmacol. Exp. Ther.*, 280(1): 67-72 (1997), respectively).

In a preferred embodiment, 5-HT₃ receptor antagonist activity of a compound can be determined according to the method described in Eguchi *et al.*, *Arzneim.-Forschung/Drug Res.*, 47(12): 1337-47 (1997). Specifically, the binding assays for determination of binding to the 5-HT₃ receptor can be performed on N1E-115 mouse neuroblastoma cells (American Type Culture Collection (ATCC) Accession No. CRL-2263) in 20 mmol/L HEPES buffer (pH=7.4) containing 150 mmol/L NaCl, 0.35 mmol/L of radiolabeled ligand ([³H]GR65630) and the test compound at 6 or more concentrations at 25 °C for 60 minutes. The reaction is terminated by rapid vacuum filtration onto glass fiber filter. Radioactivity trapped on the filter is measured by scintillation spectrometry. Non-specific binding is determined using 1 μmol/L of MDL-7222 (endo-8-methyl-8-azabicyclo [3.2.1]oct-3-yl-3,5-dichlorobenzoate. IC₅₀ values are calculated by nonlinear regression analysis. The affinity constants, K_i values, are calculated from the IC₅₀ values using the Cheng-Prusoff equation.

Compounds having 5-HT₃ receptor antagonist activity which are suitable for use in the invention have an affinity for 5-HT₃ receptor (K_i) of not more than about 250 times the K_i of ondansetron for 5-HT₃ receptor. This relative activity to ondansetron (K_i of test agent for 5-HT₃ receptor / K_i of ondansetron for 5-HT₃ receptor), can be determined by assaying the compound of interest and ondansetron using a suitable assay under controlled conditions, for example, conditions which differ primarily in the agent being tested. It is preferred that the relative activity of the 5-HT₃ receptor antagonist activity be not more than about 200 times that of ondansetron, for example, not more than about 150 times that of ondansetron, such as not more than about 100 times that of ondansetron, for example, not more than about 50 times that of ondansetron. In a particularly preferred embodiment, the

-21-

compound having 5-HT₃ receptor antagonist activity has a relative activity to ondansetron of not more than about 10.

In certain embodiments, the 5-HT₃ receptor antagonist can be selected from indisetron, YM-114 ((R)-2,3-dihydro-1-[(4,5,6,7-tetrahydro-1H-benzimidazol-5-yl)-carbonyl]-1H-indole), granisetron, talipexole, azasetron, bemasetron, tropisetron, ramosetron, ondansetron, palonosetron, lerisetron, alosetron, N-3389, zacopride, cilansetron, E-3620 ([3(S)-endo]-4-amino-5-chloro-N-(8-methyl-8-azabicyclo[3.2.1]-oct-3-yl-2[(1-methyl-2-butynyl)oxy]benzamide), lintopride, KAE-393, itasetron, zatosectron, dolasetron, (±)-zacopride, (±)-renzapride, (-)-YM-060, DAU-6236, BIMU-8 and GK-128 [2-[2-methylimidazol-1-yl)methyl]-benzo[f]thiochromen-1-one monohydrochloride hemihydrate].


In preferred embodiments, the 5-HT₃ receptor antagonist can be selected from indisetron, granisetron, azasetron, bemasetron, tropisetron, ramosetron, ondansetron, palonosetron, lerisetron, alosetron, cilansetron, itasetron, zatosectron, and dolasetron.

The invention relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need of treatment. The method comprises administering to a subject in need of treatment a therapeutically effective amount of a compound that has 5-HT₃ receptor antagonist activity and NARI activity.

In a particular embodiment, the compounds having 5-HT₃ receptor antagonist activity and NARI activity are thieno[2,3-d]pyrimidine derivatives such as those described in U.S. Patent No. 4,695,568, the entire content of which is incorporated herein by reference.

In a specific embodiment, the compounds having 5-HT₃ receptor antagonist activity and NARI activity are represented by Formula I:


-22-



wherein, R_1 and R_2 independently represent hydrogen, halogen or a C_1 - C_6 alkyl group; or R_1 and R_2 together with the carbon atoms to which they are attached form a cycloalkylene group having 5 to 6 carbon atoms;

5 R_3 and R_4 independently represent hydrogen or a C_1 - C_6 alkyl group;

R_5 is hydrogen, C_1 - C_6 alkyl,



or $-C(O)-NH-R_6$,

10 wherein m is an integer from about 1 to about 3, X is halogen and R_6 is a C_1 - C_6 alkyl group;

Ar is a substituted or unsubstituted phenyl, 2-thienyl or 3-thienyl group; and n is 2 or 3; or a pharmaceutically acceptable salt thereof.

Substituted phenyl, 2-thienyl or 3-thienyl group refers to a phenyl, 2-thienyl
 15 or 3-thienyl group in which at least one of the hydrogen atoms available for substitution has been replaced with a group other than hydrogen (i.e., a substituent group). Multiple substituent groups can be present on the phenyl, 2-thienyl or 3-thienyl ring. When multiple substituents are present, the substituents can be the same or different and substitution can be at any of the substitutable sites on the ring.
 20 Substituent groups can be, for example, a halogen atom (fluorine, chlorine, bromine

or iodine); an alkyl group, for example, a C₁-C₆ alkyl group such as a methyl, ethyl, propyl, butyl, pentyl or hexyl group; an alkoxy group, for example, a C₁-C₆ alkoxy group such as methoxy, ethoxy, propoxy, butoxy; a hydroxy group; a nitro group; an amino group; a cyano group; or an alkyl substituted amino group such as

5 methylamino, ethylamino, dimethylamino or diethylamino group.

Alkyl group refers to a straight chain or branched alkyl group. C₁-C₆ alkyl group refers to a straight-chain or branched alkyl group having from one to six carbon atoms. For example, the C₁-C₆ alkyl group can be a strain-chain alkyl such as methyl, ethyl, propyl, etc. Alternatively, the alkyl group can be branched for
10 example, an isopropyl or t-butyl group.

Halogen refers to fluorine, chlorine, bromine or iodine.

In a particular embodiment, the compounds having 5-HT₃ receptor antagonist activity and NARI activity are represented by Formula I, wherein R₁ is a C₁-C₆ alkyl group and Ar is a substituted phenyl. In this embodiment, it is preferred that the
15 phenyl group is substituted with a halogen.


In a particularly preferred embodiment, the compounds having 5-HT₃ receptor antagonist activity and NARI activity are represented by Formula I, wherein n is 2, R₁ is a C₁-C₆ alkyl group and Ar is a substituted phenyl. Preferably, the phenyl group is substituted with a halogen and the alkyl group of R₁ is a methyl
20 group.

In yet another embodiment, the compounds having 5-HT₃ receptor antagonist activity and NARI activity are represented by Formula I, wherein R₁ is a C₁-C₆ alkyl group or a halogen and Ar is an unsubstituted phenyl. Further, when R₁ is an alkyl group and Ar is an unsubstituted phenyl, R₂ can also be a hydrogen or a C₁-C₆ alkyl
25 group.

In a particularly preferred embodiment, the compounds having 5-HT₃ receptor antagonist activity and NARI activity are represented by Formula I, wherein n is 2, R₁ is a C₁-C₆ alkyl group and Ar is an unsubstituted phenyl. In a specific embodiment, wherein n is 2, R₁ is a C₁-C₆ alkyl group and Ar is an unsubstituted
30 phenyl, R₂ can be hydrogen or a C₁-C₆ alkyl group.

-24-

In a particularly preferred embodiment, the compound having 5-HT₃ receptor antagonist activity and NARI activity is represented by structural Formula II:



5 or a pharmaceutically acceptable salt thereof. This compound is referred to as MCI-225 or DDP-225. The chemical name of the structure set forth in the formula is: 4-(2-fluorophenyl)-6-methyl-2-(1-piperazinyl)thieno[2,3-d]pyrimidine.

10 In another embodiment, the method further comprises administering a therapeutically effective amount of an (i.e., one or more) additional therapeutic agent.

Compounds having 5-HT₃ receptor antagonist activity and NARI activity, such as the compounds represented by structural Formulas I and II are useful for treating nausea, vomiting, retching or any combination thereof by virtue of the dual therapeutic modes of action which they can exhibit. That is, the unique ability to
15 modulate the function of both the 5-HT₃ receptor and the noradrenaline reuptake mechanism can provide an enhanced treatment regimen for the subject undergoing treatment.

In a preferred embodiment, compounds having 5-HT₃ receptor antagonist activity and NARI activity, such as the compounds of Formula I and II possess one
20 or more characteristics selected from the group consisting of:

- a) the substantial absence of anticholinergic effects;
- b) the selective inhibition of noradrenaline reuptake as compared to inhibition of serotonin reuptake; and

- c) the selective inhibition of noradrenaline reuptake as compared to inhibition of dopamine reuptake.

For example, the specific compound MCI-225 has been shown to be a selective NARI and a 5-HT₃ receptor antagonist with substantially no anticholinergic activity. Eguchi *et al.*, *Arzneim.-Forschung/Drug Res.*, 47(12): 1337-47 (1997), reported inhibition constants for MCI-225 for the uptake the [³H]monoamine neurotransmitters noradrenaline, serotonin and dopamine in various rat brain tissues. More specifically, MCI-225 inhibited the uptake of [³H]NA and [³H]5-HT by synaptosomes from rat hypothalamic tissue with inhibition constants of Ki=35.0 nmol/L and Ki=491 nmol/L, respectively. In addition, MCI-225 inhibited the uptake of [³H]NA and [³H]5-HT by synaptosomes from rat cortical tissue with inhibition constants of Ki=0.696 nmol/L and Ki=1070 nmol/L, respectively. MCI-225 was also reported to inhibit the uptake of serotonin by synaptosomes from rat hippocampal tissue with an inhibition constant of Ki=244 nmol/L. Further, the MCI-225 inhibition constant for the uptake of [³H]DA by synaptosomes from rat striatal tissue was reported as Ki=14,800. MCI-225 did not inhibit Monoamine Oxidase-A (MAO-A) and Monoamine Oxidase-B (MAO-B) activities.

With regard to 5-HT₃ receptor antagonist activity, Eguchi *et al.* reported that MCI-225 showed high affinity for the 5-HT₃ receptor (Ki less than 100 nmol/L) in comparison to the other receptors tested. In addition, MCI-225 showed affinity for the 5-HT₃ receptor similar to that reported for ondansetron in the same radioligand binding assay. Briefly, the inhibition of radiolabeled ligand binding by MCI-225, using a suitable radioligand and tissue combination for the receptor of interest was determined. The receptors tested included, α_1 , α_2 , β_1 , β_2 , 5-HT₁, 5-HT_{1A}, 5-HT_{1c}, 5-HT₂, 5-HT₃, 5-HT₄, 5-HT₆, 5-HT₇, D₁, D₂, Muscarinic, M₁, M₂, M₃, Nicotonic, H₁, H₂, GABA-A, GABA-B, BZP, Opiate non-selective, Opiate κ , Opiate μ , Opiate δ , CRF (Corticotropin Releasing Factor) and glucocorticoid. The IC₅₀ values determined for MCI-225, for these additional receptors were all greater than 1 μ mol/L.

-26-

In one embodiment, the nausea, vomiting, retching or any combination thereof can be caused by anesthetics, radiation, cancer chemotherapeutic agents, toxic agents, odors, medicines, pregnancy and motion.

In a particular embodiment, the nausea, vomiting, retching or any
5 combination thereof can be caused by administration of general anesthetics associated with surgical procedures.

In a more particular embodiment, the nausea, vomiting, retching or any combination thereof can be caused by administration of chemotherapeutic agents, radiation therapy or a combination thereof.

10 In yet another embodiment, the nausea, vomiting, retching or any combination thereof can be caused by conditions which are associated with vertigo. For example, the nausea, vomiting, retching or any combination thereof can be caused by Meniere's disease or vestibular neuronitis.

In another embodiment, the nausea, vomiting, retching or any combination
15 thereof can be caused by headache. In particular embodiments, the headache can be a result of migraine, increased intracranial pressure or cerebral vascular hemorrhage.

In yet another embodiment, the nausea, vomiting, retching or any combination thereof can be caused by maladies of the gastrointestinal (GI) tract. In a particular embodiment, the malady of the gastrointestinal tract can be selected from
20 the group consisting of cholecystitis, choledocholithiasis, intestinal obstruction, acute gastroenteritis, perforated viscus, dyspepsia and Zollinger-Ellison syndrome.

In a further embodiment, the vomiting, nausea, retching or any combination thereof, can be of undetermined etiology. In a particular embodiment, the nausea, vomiting, retching or any combination thereof can be characterized as Cyclic
25 Vomiting Syndrome.

The invention further relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need thereof, comprising coadministering to said subject a therapeutically effective amount of a 5-HT₃ receptor antagonist and a therapeutically effective amount of a NARI.

30 The invention further relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need thereof, comprising

-27-

coadministering to said subject a first amount of a 5-HT₃ receptor antagonist and a second amount of a NARI, wherein the first and second amounts together comprise a therapeutically effective amount.

In one embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof can be caused by anesthetics, radiation, cancer chemotherapeutic agents, toxic agents, odors, medicines, pregnancy and motion.

In a particular embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof can be caused by administration of general anesthetics associated with surgical procedures.

In a more particular embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof caused by administration of chemotherapeutic agents, radiation therapy or a combination thereof.

In yet another embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof caused by conditions which are associated with vertigo. For example, the nausea, vomiting, retching or any combination thereof can be caused by Meniere's disease or vestibular neuronitis.

In another embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof caused by headache. In particular embodiments, the headache is a result of migraine, increased intracranial pressure or cerebral vascular hemorrhage.

In yet another embodiment, the coadministration can be used to treat nausea, vomiting, retching or any combination thereof caused by maladies of the gastrointestinal (GI) tract. In a particular embodiment, the malady of the gastrointestinal tract is selected from the group consisting of cholecystitis, choledocholithiasis, intestinal obstruction, acute gastroenteritis, perforated viscus, dyspepsia and Zollinger-Ellison syndrome.

In a further embodiment, the coadministration can be used to treat vomiting, nausea, retching or any combination thereof, of undetermined etiology. In a particular embodiment, the nausea, vomiting, retching or any combination thereof can be characterized as Cyclic Vomiting Syndrome.

In another embodiment, the coadministration methods further comprise administering a therapeutically effective amount of an (i.e., one or more) additional therapeutic agent.

In certain embodiments of the coadministration method, the 5-HT₃ receptor antagonist can be selected from indisetron, YM-114 ((R)-2,3-dihydro-1-[(4,5,6,7-tetrahydro-1H-benzimidazol-5-yl)-carbonyl]-1H-indole), granisetron, talipexole, azasetron, bemesetron, tropisetron, ramosetron, ondansetron, palonosetron, lerisetron, alosetron, N-3389, zacopride, cilansetron, E-3620 ([3(S)-endo]-4-amino-5-chloro-N-(8-methyl-8-azabicyclo[3.2.1]-oct-3-yl-2[(1-methyl-2-butynyl)oxy]benzamide), lintopride, KAE-393, itasetron, zatosetron, dolasetron, (±)-zacopride, (±)-renzapride, (-)-YM-060, DAU-6236, BIMU-8 and GK-128 [2-[2-methylimidazol-1-yl)methyl]-benzo[f]thiochromen-1-one monohydrochloride hemihydrate].

In preferred embodiments, the 5-HT₃ receptor antagonist can be selected from indisetron, granisetron, azasetron, bemesetron, tropisetron, ramosetron, ondansetron, palonosetron, lerisetron, alosetron, cilansetron, itasetron, zatosetron, and dolasetron.

In certain embodiments, the NARI compound can be selected from venlafaxine, duloxetine, bupropion, milnacipran, reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotiline, viloxazine and atomoxetine.

In a preferred embodiment, the NARI compound can be selected from reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotiline, viloxazine and atomoxetine.

In a preferred embodiment, the NARI compound possesses one or more characteristics selected from the group consisting of:

- a) the substantial absence of anticholinergic effects;
- b) the selective inhibition of noradrenaline reuptake as compared to inhibition of serotonin reuptake; and
- c) the selective inhibition of noradrenaline reuptake as compared to inhibition of dopamine reuptake.

In addition, the invention relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need thereof comprising administering a therapeutically effective amount of a NARI. In this embodiment, the NARI is characterized by the substantial absence of anticholinergic effects.

5 In a further embodiment, the NARI possesses selective inhibition of noradrenaline reuptake as compared to inhibition of serotonin reuptake and/or selective inhibition of noradrenaline reuptake as compared to inhibition of dopamine reuptake.

10 In one embodiment, the nausea, vomiting, retching or any combination thereof can be caused by anesthetics, radiation, cancer chemotherapeutic agents, toxic agents, odors, medicines, pregnancy and motion.

In a particular embodiment, the nausea, vomiting, retching or any combination thereof can be caused by administration of general anesthetics associated with surgical procedures.

15 In a more particular embodiment, the nausea, vomiting, retching or any combination thereof can be caused by administration of chemotherapeutic agents, radiation therapy or a combination thereof.

20 In yet another embodiment, the nausea, vomiting, retching or any combination thereof can be caused by conditions which are associated with vertigo. For example, the nausea, vomiting, retching or any combination thereof can be caused by Meniere's disease or vestibular neuronitis.

In another embodiment, the nausea, vomiting, retching or any combination thereof can be caused by headache. In particular embodiments, the headache can be a result of migraine, increased intracranial pressure or cerebral vascular hemorrhage.

25 In yet another embodiment, the nausea, vomiting, retching or any combination thereof can be caused by maladies of the gastrointestinal (GI) tract. In a particular embodiment, the malady of the gastrointestinal tract can be selected from the group consisting of cholecystitis, choledocholithiasis, intestinal obstruction, acute gastroenteritis, perforated viscus, dyspepsia and Zollinger-Ellison syndrome.

In a further embodiment, the vomiting, nausea, retching or any combination thereof, is of undetermined etiology. In a particular embodiment, the nausea, vomiting, retching or any combination thereof is Cyclic Vomiting Syndrome.

In another embodiment, the method further comprises administering a
5 therapeutically effective amount of an (i.e., one or more) additional therapeutic agent. For example, the method can further comprise administering a corticosteroid.

The invention further relates to pharmaceutical compositions useful for the treatment of a nausea, vomiting, retching or any combination thereof. The pharmaceutical composition comprises a first amount of a 5-HT₃ receptor antagonist
10 compound and a second amount of a NARI compound. The pharmaceutical compositions of the present invention can optionally contain a pharmaceutically acceptable carrier. The 5-HT₃ receptor antagonist and the NARI can each be present in the pharmaceutical composition in a therapeutically effective amount. In another aspect, said first and second amounts can together comprise a therapeutically
15 effective amount.

In a further embodiment, the pharmaceutical composition further comprises an (i.e., one or more) additional therapeutic agent.

The pharmaceutical composition can be used in the treatment of a nausea, vomiting, retching or any combination thereof in a subject in need of treatment. As
20 such, the invention relates to a method of treating nausea, vomiting, retching or any combination thereof in a subject in need of treatment comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition as described herein.

An additional therapeutic agent suitable for use in the methods and
25 pharmaceutical compositions described herein, can be, but is not limited to, for example: an anticholinergic (e.g., scopolomine); an antihistamine (e.g., dimenhydrinate and diphenhydramine); a phenothiazine (e.g., prochlorperazine and chlorpromazine); a butyrophenone (haloperidol and droperidol); a cannabinoid (e.g., tetrahydrocannabinol and nabilone); a benzamide (e.g., metoclopramide, cisapride
30 and trimethobenzamide); a glucocorticoid (e.g., dexamethasone and methylprednisolone); a benzodiazepine (e.g., lorazepam); or any combination

-31-

thereof. In a preferred embodiment, the additional therapeutic agent is a glucocorticoid.

VOMITING, NAUSEA AND RETCHING

Emesis and vomiting, as used herein, are synonymous and can be described
5 as the forceful expulsion of gastrointestinal contents through the mouth brought about by the descent of the diaphragm and powerful contractions of the abdominal muscles. Emesis is usually, but not always, preceded by nausea.

Nausea, as used herein, is the unpleasant feeling that one is about to vomit.

Retching or dry heaves, as used herein, involves the same physiological
10 mechanisms as vomiting, but occurs against a closed glottis, which prohibits the expulsion of gastric contents.

Nausea, vomiting, retching or combinations thereof can be caused by a number of factors including, but not limited to, anesthetics, radiation, cancer
chemotherapeutic agents, toxic agents, odors, medicines, for example, serotonin
15 reuptake inhibitors, analgesics such as morphine, antibiotics and antiparasitic agents, pregnancy and motion. Conditions which are associated with vertigo (e.g., Meniere's disease and vestibular neuronitis) can also cause nausea, vomiting, retching or any combination thereof. Headache, caused by, for example, migraine, increased intracranial pressure or cerebral vascular hemorrhage can also result in
20 nausea, vomiting, retching or any combination thereof. In addition, certain maladies of the gastrointestinal (GI) tract, for example, cholecystitis, choledocholithiasis, intestinal obstruction, acute gastroenteritis, perforated viscus, dyspepsia resulting from, for example, gastroesophageal reflux disease, peptic ulcer disease, gastroparesis, gastric or esophageal neoplasms, infiltrative gastric disorders (e.g.,
25 Menetrier's syndrome, Crohn's disease, eosinophilic gastroenteritis, sarcoidosis and amyloidosis) gastric infections (e.g., CMV, fungal, TB and syphilis), parasites (e.g., *Giardia lamblia* and *Strongyloides stercoralis*), chronic gastric volvulus, chronic intestinal ischemia, altered gastric motility and/or food intolerance or Zollinger-Ellison syndrome can result in vomiting, nausea, retching or any combination
30 thereof. However, in some cases of vomiting, nausea, retching or combinations

-32-

thereof, no etiology can be determined despite extensive diagnostic testing(e.g., Cyclic Vomiting Syndrome).

Nausea, vomiting and retching are defined as acute when symptoms are present for less than a week. The causes of nausea, vomiting and retching of short
5 duration are often separable from etiologies leading to more chronic symptoms.

Nausea, vomiting and retching are defined as chronic when symptoms are present for over a week. For example, symptoms can be continuous or intermittent and last for months or years.

Dyspepsia, as used herein, refers to pain or discomfort centered in the upper
10 abdomen that can also include bloating, early satiety, postprandial fullness, nausea, anorexia, heartburn, regurgitation, and burping or belching. Generally, the symptoms of dyspepsia arise from the upper luminal GI tract. Dyspepsia can be caused by a number of foods, medications, systemic disorders and diseases of the luminal GI tract.

15 Subject, as used herein, refers to animals such as mammals, including, but not limited to, primates (e.g., humans), cows, sheep, goats, horses, pigs, dogs, cats, rabbits, guinea pigs, rats, mice or other bovine, ovine, equine, canine, feline, rodent or murine species.


As used herein, therapeutically effective amount refers to an amount
20 sufficient to elicit the desired biological response. In the present invention the desired biological response is a reduction (complete or partial) in vomiting, nausea, retching or any combination thereof resulting from any cause.

CHEMOTHERAPEUTIC AGENTS

Chemotherapeutic agents, as that term is used herein, include, but are not
25 limited to, for example alkylating agents, e.g. cyclophosphamide, carmustine, lomustine, and chlorambucil; cytotoxic antibiotics, e.g. dactinomycin, doxorubicin, mitomycin-C, and bleomycin; antimetabolites, e.g. cytarabine, methotrexate, and 5-fluorouracil; vinca alkaloids, e.g. etoposide, vinblastine, and vincristine; and others such as cisplatin, dacarbazine, procarbazine, and hydroxyurea; and
30 combinations thereof.

MEDICINES

In addition to chemotherapeutic agents, many other medicines can cause nausea, vomiting, retching or a combination thereof. For example, analgesics, antibiotics, antiparasitic agents, and serotonin reuptake inhibitors can cause nausea, vomiting, retching or a combination thereof. A serotonin reuptake inhibitor (SRI) is any compound which inhibits the uptake of serotonin. In a particular embodiment, the nausea, vomiting, retching or any combination thereof is associated with the onset of SRI therapy. When the SRI is dosed on an as needed basis (prn), each dose can be considered the onset of therapy and can cause nausea, vomiting, retching or any combination thereof. Such prn dosing is typically used in the treatment of premature ejaculation. The SRI can have other therapeutic characteristics such as inhibition of the uptake of noradrenaline. Therefore, SRIs include selective serotonin reuptake inhibitors (SSRIs) such as fluoxetine, paroxetine, sertraline and the rapid onset SSRI dapoxetine. In addition, certain SSRIs are known to exhibit 5-HT_{1A} receptor activities (e.g., antagonist or partial agonist activity at the 5-HT_{1A} receptor). Compounds which have combined SSRI and 5-HT_{1A} receptor activities include those described in WO 99/02516 and WO 02/44170, the contents of which are incorporated herein by reference. These compounds are represented by the Formulas III, IV and V:



wherein, Z is -CO-, -CH(OR₆)- or -C(NOR₇)-;

R₁ is hydrogen, C₁-C₆ alkyl, halogen or -O-R₁₂;

R₂ and R₃ independently represent hydrogen, C₁-C₆ alkyl, halogen, nitro or -O-R₆

25 or R₂ and R₃ are together -CR₈=CR₉-CR₁₀=CR₁₁;

-34-

R₄ and R₅ independently represent hydrogen, alkyl, halogen, haloalkyl, -OR₁₂, nitro, -NR₁₃R₁₄, -COR₁₂, -CO₂R₁₂, -SO₂NR₁₃R₁₄, -SO₂R₁₂, -SR₁₂, cyano, -CONR₁₃R₁₄;

or R₄ and R₅ together form a benzene ring;

5 R₆ is hydrogen, C₁-C₆ alkyl, -CO₂R₁₂, -C(O)NR₁₃R₁₄, naphthyl or phenyl;


R₇ is hydrogen or C₁-C₆ alkyl;

R₈, R₉, R₁₀ and R₁₁ are independently hydrogen, C₁-C₆ alkyl, halogen, -OR₁₂, nitro, cyano, -NR₁₃R₁₄, -COR₁₂, -CO₂R₁₂, SO₂NR₁₃R₁₄, -SO₂R₁₂, -SR₁₂, or -CONR₁₃R₁₄;


10 R₁₂ is hydrogen, C₁-C₆ alkyl or phenyl; and

R₁₃ and R₁₄ are independently hydrogen, C₁-C₆ alkyl or phenyl or

R₁₃ and R₁₄ form a ring of 5 or 6 members; or a pharmaceutically acceptable salt or solvate or any isomer (geometric or optical) or polymorph thereof.



15 wherein, the variables Z, R₁, R₂, R₃, R₄ and R₅ of Formula IV have the meanings set forth in Formula III.



-35-

wherein, n is 1, 2 or 3;

Z is -C(O) or -CHOH;

R₁ is hydrogen, C₁-C₆ alkyl, halogen -OR₂, nitro, cyano, -NR₃R₄, -COR₂,
-CO₂R₂, -O-COR₂, -SO₂NR₃R₄, -SO₂R₂, -SR₂ or -CONR₃R₄;

5 R₂ is hydrogen, C₁-C₆ alkyl or phenyl;

R₃ and R₄ are independently hydrogen, C₁-C₆ alkyl or phenyl or

R₃ and R₄ together form a morpholine, thiomorphone or piperazine ring;

Ar is an optionally substituted bicyclic system formed by a benzocondensed
heterocyclic ring with 5, 6 or 7 ring atoms, saturated or unsaturated and

10 containing 1, 2 or 3 heteroatoms selected from N, O or S; or a

pharmaceutically acceptable salt or solvate or any isomer (geometric or
optical) or polymorph thereof.

MODES OF ADMINISTRATION

The compounds for use in the method of the invention can be formulated for
15 oral, transdermal, sublingual, buccal, parenteral, rectal, intranasal, intrabronchial or
intrapulmonary administration. For oral administration the compounds can be of the
form of tablets or capsules prepared by conventional means with pharmaceutically
acceptable excipients such as binding agents (e.g., polyvinylpyrrolidone,
hydroxypropylcellulose or hydroxypropylmethylcellulose); fillers (e.g., cornstarch,
20 lactose, microcrystalline cellulose or calcium phosphate); lubricants (e.g.,
magnesium stearate, talc, or silica); disintegrates (e.g., sodium starch glycollate); or
wetting agents (e.g., sodium lauryl sulphate). If desired, the tablets can be coated
using suitable methods and coating materials such as OPADRY® film coating
systems available from Colorcon, West Point, PA (e.g., OPADRY® OY Type, OY-C
25 Type, Organic Enteric OY-P Type, Aqueous Enteric OY-A Type, OY-PM Type and
OPADRY® White, 32K18400). Liquid preparation for oral administration can be in
the form of solutions, syrups or suspensions. The liquid preparations can be
prepared by conventional means with pharmaceutically acceptable additives such as
suspending agents (e.g., sorbitol syrup, methyl cellulose or hydrogenated edible

fats); emulsifying agent (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters or ethyl alcohol); and preservatives (e.g., methyl or propyl p-hydroxy benzoates or sorbic acid).

For buccal administration, the compounds for use in the method of the invention can be in the form of tablets or lozenges formulated in a conventional manner.

For parenteral administration, the compounds for use in the method of the invention can be formulated for injection or infusion, for example, intravenous, intramuscular or subcutaneous injection or infusion, or for administration in a bolus dose and/or continuous infusion. Suspensions, solutions or emulsions in an oily or aqueous vehicle, optionally containing other formulatory agents such as suspending, stabilizing and/or dispersing agents can be used.

For rectal administration, the compounds for use in the method of the invention can be in the form of suppositories.

For sublingual administration, tablets can be formulated in conventional manner.

For intranasal, intrabronchial or intrapulmonary administration, conventional formulations can be employed.

Further, the compounds for use in the method of the invention can be formulated in a sustained release preparation. For example, the compounds can be formulated with a suitable polymer or hydrophobic material which provides sustained and/or controlled release properties to the active agent compound. As such, the compounds for use the method of the invention can be administered in the form of microparticles for example, by injection or in the form of wafers or discs by implantation.

Additional dosage forms of this invention include dosage forms as described in U.S. Pat. No. 6,340,475, U.S. Pat. No. 6,488,962, U.S. Pat. No. 6,451,808, U.S. Pat. No. 6,340,475, U.S. Pat. No. 5,972,389, U.S. Pat. No. 5,582,837, and U.S. Pat. No. 5,007,790. Additional dosage forms of this invention also include dosage forms as described in U.S. Pat. Application No. 20030147952, U.S. Pat. Application No. 20030104062, U.S. Pat. Application No. 20030104053, U.S. Pat. Application No.

20030044466, U.S. Pat. Application No. 20030039688, and U.S. Pat. Application No. 20020051820. Additional dosage forms of this invention also include dosage forms as described in PCT Patent Application WO 03/35041, PCT Patent Application WO 03/35040, PCT Patent Application WO 03/35029, PCT Patent Application WO 03/35177, PCT Patent Application WO 03/35039, PCT Patent Application WO 02/96404, PCT Patent Application WO 02/32416, PCT Patent Application WO 01/97783, PCT Patent Application WO 01/56544, PCT Patent Application WO 01/32217, PCT Patent Application WO 98/55107, PCT Patent Application WO 98/11879, PCT Patent Application WO 97/47285, PCT Patent Application WO 93/18755, and PCT Patent Application WO 90/11757.

In one embodiment, the dosage forms of the present invention include pharmaceutical tablets for oral administration as described in U.S. Patent Application No. 20030104053. For example, suitable dosage forms of the present invention can combine both immediate-release and prolonged-release modes of drug delivery. The dosage forms of this invention include dosage forms in which the same drug is used in both the immediate-release and the prolonged-release portions as well as those in which one drug is formulated for immediate release and another drug, different from the first, is formulated for prolonged release. This invention encompasses dosage forms in which the immediate-release drug is at most sparingly soluble in water, i.e., either sparingly soluble or insoluble in water, while the prolonged-release drug can be of any level of solubility.

More particularly, in a further embodiment, the prolonged-release portion of the dosage form can be a dosage form that delivers its drug to the digestive system continuously over a period of time of at least an hour and preferably several hours and the drug is formulated as described in U.S. Patent Application No. 20030104053. In said embodiment, the immediate-release portion of the dosage form can be a coating applied or deposited over the entire surface of a unitary prolonged-release core, or can be a single layer of a tablet constructed in two or more layers, one of the other layers of which is the prolonged-released portion and is formulated as described in U.S. Patent Application No. 20030104053.

-38-

In another embodiment of the invention, the supporting matrix in controlled-release tablets or controlled release portions of tablets is a material that swells upon contact with gastric fluid to a size that is large enough to promote retention in the stomach while the subject is in the digestive state, which is also referred to as the postprandial or "fed" mode. This is one of two modes of activity of the stomach that differ by their distinctive patterns of gastroduodenal motor activity. The "fed" mode is induced by food ingestion and begins with a rapid and profound change in the motor pattern of the upper gastrointestinal (GI) tract. The change consists of a reduction in the amplitude of the contractions that the stomach undergoes and a reduction in the pyloric opening to a partially closed state. The result is a sieving process that allows liquids and small particles to pass through the partially open pylorus while indigestible particles that are larger than the pylorus are retropelled and retained in the stomach. This process causes the stomach to retain particles that are greater than about 1 cm in size for about 4 to 6 hours. The controlled-release matrix in these embodiments of the invention is therefore selected as one that swells to a size large enough to be retropelled and thereby retained in the stomach, causing the prolonged release of the drug to occur in the stomach rather than in the intestines. Disclosures of oral dosage forms that swell to sizes that will prolong the residence time in the stomach are found in U.S. Pat. No. 6,448,962, U.S. Pat. No. 6,340,475, U.S. Pat. No. 5,007,790, U.S. Pat. No. 5,582,837, U.S. Pat. No. 5,972,389, PCT Patent Application WO 98/55107, U.S. Patent Application No. 20010018707, U.S. Patent Application No. 20020051820, U.S. Patent Application No. 20030029688, U.S. Patent Application No. 20030044466, U.S. Patent Application No. 20030104062, U.S. Patent Application No. 20030147952, U.S. Patent Application No. 20030104053, and PCT Patent Application WO 96/26718. In particular, gastric retained dosage formulations for specific drugs have also been described, for example, a gastric retained dosage formulation for gabapentin is disclosed in PCT Patent Application WO 03/035040.

COADMINISTRATION

In practicing the methods of the invention, coadministration refers to administration of a first amount of a 5-HT₃ receptor antagonist compound and a second amount of a NARI compound to treat nausea, vomiting, retching or any combination thereof. Coadministration encompasses administration of the first and second amounts of the compounds of the coadministration in an essentially simultaneous manner, such as in a single pharmaceutical composition, for example, capsule or tablet having a fixed ratio of first and second amounts, or in multiple, separate capsules or tablets for each. In addition, such coadministration also encompasses use of each compound in a sequential manner in either order. When coadministration involves the separate administration of the NARI and 5-HT₃ receptor antagonist, the compounds are administered sufficiently close in time to have the desired therapeutic effect.

DOSING

The therapeutically effective amount or dose of (a) a compound having dual therapeutic modes of action (i.e., 5-HT₃ receptor antagonist activity and NARI activity); (b) a 5-HT₃ receptor antagonist and NARI in combination; or (c) a NARI alone, will depend on the age, sex and weight of the patient, the current medical condition of the patient and the nature of the nausea, vomiting, retching or any combination thereof being treated. The skilled artisan will be able to determine appropriate dosages depending on these and other factors.

As used herein, continuous dosing refers to the chronic administration of a selected active agent.

As used herein, as-needed dosing, also known as "pro re nata" "prn" dosing, and "on demand" dosing or administration is meant the administration of a therapeutically effective dose of the compound(s) at some time prior to commencement of an activity wherein suppression of nausea, vomiting, retching or any combination thereof would be desirable. Administration can be immediately prior to such an activity, including about 0 minutes, about 10 minutes, about 20 minutes, about 30 minutes, about 1 hour, about 2 hours, about 3 hours, about 4

hours, about 5 hours, about 6 hours, about 7 hours, about 8 hours, about 9 hours, or about 10 hours prior to such an activity, depending on the formulation.

In a particular embodiment, drug administration or dosing is on an as-needed basis, and does not involve chronic drug administration. With an immediate release dosage form, as-needed administration can involve drug administration immediately prior to commencement of an activity wherein suppression of nausea, vomiting, retching or any combination thereof would be desirable, but will generally be in the range of from about 0 minutes to about 10 hours prior to such an activity, preferably in the range of from about 0 minutes to about 5 hours prior to such an activity, most preferably in the range of from about 0 minutes to about 3 hours prior to such an activity.

For example, a suitable dose of the 5-HT₃ receptor antagonist can be in the range of from about 0.001 mg to about 500 mg per day, such as from about 0.01 mg to about 100 mg, for example, from about 0.05 mg to about 50 mg, such as about 0.5 mg to about 25 mg per day. The dose can be administered in a single dosage or in multiple dosages, for example from 1 to 4 or more times per day. When multiple dosages are used, the amount of each dosage can be the same or different.

For example, a suitable dose of the NARI compound can be in the range of from about 0.001 mg to about 1000 mg per day, such as from about 0.05 mg to about 500 mg, for example, from about 0.03 mg to about 300 mg, such as about 0.02 mg to about 200 mg per day. The dose can be administered in a single dosage or in multiple dosages, for example from 1 to 4 or more times per day. When multiple dosages are used, the amount of each dosage can be the same or different.

For example, a suitable dose of the compound having both 5-HT₃ receptor antagonist and NARI activity can be in the range of from about 0.001 mg to about 1000 mg per day, such as from about 0.05 mg to about 500 mg, for example, from about 0.03 mg to about 300 mg, such as from about 0.02 mg to about 200 mg per day. In a particular embodiment, a suitable dose of the compound having both 5-HT₃ receptor antagonist and NARI activity can be in the range of from about 0.1 mg to about 50 mg per day, such as from about 0.5 mg to about 10 mg per day such as about 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 mg per day. The dose per day can be

administered in a single dosage or in multiple dosages, for example from 1 to 4 or more times per day. When multiple dosages are used, the amount of each dosage can be the same or different. For example a dose of 1 mg per day can be administered as two 0.5 mg doses, with about a 12 hour interval between doses.

5 It is understood that the amount of compound dosed per day can be administered every day, every other day, every 2 days, every 3 days, every 4 days, every 5 days, etc. For example, with every other day administration, a 5 mg per day dose can be initiated on Monday with a first subsequent 5 mg per day dose administered on Wednesday, a second subsequent 5 mg per day dose administered
10 on Friday, etc.

 The compounds for use in the method of the invention can be formulated in unit dosage form. The term "unit dosage form" refers to physically discrete units suitable as unitary dosage for subjects undergoing treatment, with each unit containing a predetermined quantity of active material calculated to produce the
15 desired therapeutic effect, optionally in association with a suitable pharmaceutical carrier. The unit dosage form can be for a single daily dose or one of multiple daily doses (e.g., about 1 to 4 or more times per day). When multiple daily doses are used, the unit dosage form can be the same or different for each dose.

 For the compounds having both NARI and 5-HT₂ receptor antagonist
20 activity, each dosage can typically contain from about 0.001 mg to about 1000 mg, such as from about 0.05 mg to about 500 mg, for example, from about 0.03 mg to about 300 mg, such as about 0.02 mg to about 200 mg of the active ingredient.

 The compounds for use in the method of the invention can be formulated in unit dosage form. The term "unit dosage form" refers to physically discrete units
25 suitable as unitary dosage for subjects undergoing treatment, with each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, optionally in association with a suitable pharmaceutical carrier. The unit dosage form can be for a single daily dose or one of multiple daily doses (e.g., about 1 to 4 or more times per day). When multiple daily doses are
30 used, the unit dosage form can be the same or different for each dose. Dosing can be on demand by the subject.

For the compounds having both NARI and 5-HT₃ receptor antagonist activity, each dosage can typically contain from about 0.001 mg to about 1000 mg, such as from about 0.05 mg to about 500 mg, for example, from about 0.03 mg to about 300 mg, such as about 0.02 mg to about 200 mg of the active ingredient.

5 When the method of treatment comprises coadministration of a NARI and a 5-HT₃ receptor antagonist each dose can typically contain from about .001 mg to about 1000 mg, such as from about 0.05 mg to about 500 mg, for example, from about 0.03 mg to about 300 mg, such as about 0.02 mg to about to about 200 mg of the NARI and typically can contain from about 0.001 mg to about 500 mg, such as
10 from about 0.01 mg to about 100 mg, for example, from about 0.05 mg to about 50 mg, such as about 0.5 mg to about 25 mg of the 5-HT₃ receptor antagonist.

 When the method of treatment comprises administration of a NARI alone, each dose can typically contain from about .001 mg to about 1000 mg, such as from about 0.05 mg to about 500 mg, for example, from about 0.03 mg to about 300 mg,
15 such as 0.02 to about to about 200 mg of the active ingredient.

 The invention further includes a kit for treating nausea, vomiting, retching or any combination thereof. The kit comprises at least one compound having both 5-HT₃ receptor antagonist activity and NARI activity (e.g., a single compound) and an instruction insert for administering the compound according to the method of the
20 invention. In addition, the kit can comprise a first compound which is a 5-HT₃ receptor antagonist and a second compound which is a NARI and an instruction insert for administering the compounds according to the method of the invention. The first and second compounds can be in separate dosage forms or combined in a single dosage form.

25 As used herein, the term pharmaceutically acceptable salt refers to a salt of the administered compounds prepared from pharmaceutically acceptable non-toxic acids including inorganic acids, organic acids, solvates, hydrates, or clathrates thereof. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, and phosphoric. Appropriate organic acids may be
30 selected, for example, from aliphatic, aromatic, carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic,

camphorsulfonic, citric, fumaric, gluconic, isethionic, lactic, malic, mucic, tartaric, para-toluenesulfonic, glycolic, glucuronic, maleic, furoic, glutamic, benzoic, anthranilic, salicylic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, pantothenic, benzenesulfonic (besylate), stearic, sulfanilic, alginic, galacturonic, and the like.

It is understood that 5-HT₃ receptor antagonists, NARIs and single compounds having both NARI and 5-HT₃ antagonist activities can be identified, for example, by screening libraries or collections of molecules using suitable methods. Another source for the compounds of interest are combinatorial libraries which can comprise many structurally distinct molecular species. Combinatorial libraries can be used to identify lead compounds or to optimize a previously identified lead. Such libraries can be manufactured by well-known methods of combinatorial chemistry and screened by suitable methods.

The invention also relates to a method of processing a claim under a health insurance policy submitted by a claimant seeking reimbursement for costs associated with the treatment of nausea, vomiting, retching or any combination thereof, as described herein.

In one embodiment, the method of processing a claim under a health insurance policy submitted by a claimant seeking reimbursement for costs associated with treatment of nausea, vomiting, retching or any combination thereof, wherein said treatment comprises coadministering to a subject a first amount of a 5-HT₃ receptor antagonist and a second amount of a noradrenaline reuptake inhibitor, wherein the first and second amounts together comprise a therapeutically effective amount comprising: reviewing said claim; determining whether said treatment is reimbursable under said insurance policy; and processing said claim to provide partial or complete reimbursement of said costs.

In one embodiment, the nausea, vomiting, retching or any combination thereof is caused by an anesthetic, radiation, a cancer chemotherapeutic agent, a toxic agent, an odor, a medicine, pregnancy or motion.

In a particular embodiment, the medicine is selected from the group consisting of an analgesic, an antibiotic, an antifungal or a serotonin reuptake inhibitor.

In another embodiment, the nausea, vomiting, retching or any combination thereof is caused by a condition which is associated with vertigo.

In another embodiment, the nausea, vomiting, retching or any combination thereof is caused by headache.

In yet another embodiment, the nausea, vomiting, retching or any combination thereof is caused by a malady of the gastrointestinal (GI) tract.

10 The invention also relates to a method for processing a claim under a health insurance policy submitted by a claimant seeking reimbursement for costs associated with treatment of nausea, vomiting, retching or any combination thereof, wherein said treatment comprises coadministering to a subject a therapeutically effective amount of a 5-HT₃ receptor antagonist and a therapeutically effective
15 amount of a noradrenaline reuptake inhibitor comprising: reviewing said claim; determining whether said treatment is reimbursable under said insurance policy; and processing said claim to provide partial or complete reimbursement of said costs.

In one embodiment, the nausea, vomiting, retching or any combination thereof is caused by an anesthetic, radiation, a cancer chemotherapeutic agent, a
20 toxic agent, an odor, a medicine, pregnancy or motion.

In a particular embodiment, the medicine is selected from the group consisting of an analgesic, an antibiotic, an antifungal or a serotonin reuptake inhibitor.

In another embodiment, the nausea, vomiting, retching or any combination thereof is caused by a condition which is associated with vertigo.

In another embodiment, the nausea, vomiting, retching or any combination thereof is caused by headache.

In yet another embodiment, the nausea, vomiting, retching or any combination thereof is caused by a malady of the gastrointestinal (GI) tract.

30 The invention also relates to a method for processing a claim under a health insurance policy submitted by a claimant seeking reimbursement for costs

associated with treatment of nausea, vomiting, retching or any combination thereof, wherein said treatment comprises administering to a subject a therapeutically effective amount of a compound having 5-HT₃ receptor antagonist activity and noradrenaline reuptake inhibitor activity comprising: reviewing said claim;
5 determining whether said treatment is reimbursable under said insurance policy; and processing said claim to provide partial or complete reimbursement of said costs.

In a particular embodiment, the compound having 5-HT₃ receptor antagonist activity and noradrenaline reuptake inhibitor activity is MCI-225.

In one embodiment, the nausea, vomiting, retching or any combination
10 thereof is caused by an anesthetic, radiation, a cancer chemotherapeutic agent, a toxic agent, an odor, a medicine, pregnancy or motion.

In a particular embodiment, the medicine is selected from the group consisting of an analgesic, an antibiotic, an antifungal or a serotonin reuptake inhibitor.

15 In another embodiment, the nausea, vomiting, retching or any combination thereof is caused by a condition which is associated with vertigo.

In another embodiment, the nausea, vomiting, retching or any combination thereof is caused by headache.

In yet another embodiment, the nausea, vomiting, retching or any
20 combination thereof is caused by a malady of the gastrointestinal (GI) tract.

The invention further relates to a method for processing a claim under a health insurance policy submitted by a claimant seeking reimbursement for costs associated with treatment of nausea, vomiting, retching or any combination thereof, wherein said treatment comprises administering to a subject a therapeutically
25 effective amount of a noradrenaline reuptake inhibitor, wherein the noradrenaline reuptake inhibitor characterized by the substantial absence of anticholinergic effects comprising: reviewing said claim; determining whether said treatment is reimbursable under said insurance policy; and processing said claim to provide partial or complete reimbursement of said costs.

-46-

In one embodiment, the nausea, vomiting, retching or any combination thereof is caused by an anesthetic, radiation, a cancer chemotherapeutic agent, a toxic agent, an odor, a medicine, pregnancy or motion.

In a particular embodiment, the medicine is selected from the group
5 consisting of an analgesic, an antibiotic, an antifungal or a serotonin reuptake inhibitor.

In another embodiment, the nausea, vomiting, retching or any combination thereof is caused by a condition which is associated with vertigo.

In another embodiment, the nausea, vomiting, retching or any combination
10 thereof is caused by headache.

In yet another embodiment, the nausea, vomiting, retching or any combination thereof is caused by a malady of the gastrointestinal (GI) tract.

PHARMACOLOGICAL METHODS

ANTI-EMETIC EFFECTS

15 The activity of compounds as anti-emetics can be demonstrated by any suitable model. For example, the extent to which compounds can reduce the latency or the number of retches and/or vomits induced by emetogens (e.g., cisplatin which is a typically used emetogenic trigger in suitable animal models) in, for example, the dog (e.g., beagles), the piglet or in the ferret can be assessed. For example, suitable
20 methods are described in Tattersall *et al.* and Bountra *et al.*, *European Journal of Pharmacology*, 250: (1993) R5 and 249:(1993) R3-R4 and Milano *et al.*, *J. Pharmacol. Exp. Ther.*, 274(2): 951-961 (1995).

In addition, the general method described by Florezyk *et al.*, *Cancer Treatment Report*, 66(1): 187-9, (1982)) and summarized below, can also be used to
25 assess effect of a test compound on emesis in the ferret.

Briefly, both the test compound and cisplatin are prepared and administered. The cisplatin is a representative emetogenic trigger for vomiting.

-47-

a) Control--Without Test Agent

Emesis is induced in groups of 6 male ferrets weighing about 2 kg, by intravenous administration of cisplatin at a suitable dose (e.g., 10 mg/kg). The onset of emesis is noted. Over a period of 2 hours the number of vomits/retches (episodes) is recorded. Behavioral changes characteristic of emesis are also noted.

b) With Test Compound

The test compound is administered to groups of 6 male ferrets weighing about 2 kg, by intravenous administration at suitable doses immediately prior to administration of cisplatin as described above. The animals are observed for 3 hours.

The emetic response seen in drug tested and control animals can then be compared to assess antiemetic properties of the test compound.

EXEMPLIFICATION

The present invention will now be illustrated by the following Example, which is not intended to be limiting in any way.

TREATMENT OF VOMITING AND RETCHING USING MCI-225

The ability of MCI-225 to reduce retching and vomiting in an accepted model of cytotoxin-induced emesis in the ferret was assessed. Specifically, the experiments described herein investigated the effect of MCI-225 on retching and vomiting induced by cisplatin. Ondansetron was used as a positive control in the model, in view of its known antiemetic activity.

Animals

Adult male ferrets (*Mustela putario furo*) weighing 1200-1880g were purchased from Triple F Farms (Sayre, PA) and housed in individual cages at standardized conditions (12:12 h light/dark cycle and 21-23°C). Prior to the experiments, the ferrets were allowed a 7-10 day acclimatization period to the

animal facility. The ferrets were fed a carnivore diet with free access to food and water throughout the course of the study. The use of the ferret model of emesis and the drug treatment were preapproved in accordance with facility standards.

Cisplatin-Induced Emesis

- 5 A cisplatin solution was prepared by adding preheated (70°C) saline to cisplatin powder (Sigma-Aldrich Co.) and stirring or sonicating at 40°C until dissolved.

- Following administration of the cisplatin and either MCI-225, ondansetron or vehicle alone, the occurrence of retching and vomiting was monitored for a
10 period of 6 hours. Retching was defined as the number of forceful rhythmic contractions of the abdomen occurring with the animal in characteristic posture, but not resulting in the expulsion of upper gastrointestinal tract contents (Watson *et al.*, *British Journal of Pharmacology*, 115(1): 84-94 (1994)). Vomiting was defined as the forceful oral expulsion of upper gastrointestinal contents. The latency of the
15 retching or vomiting response and the number of episodes were recorded for each animal and summarized for each experimental group (Wright *et al.*, *Infect. Immun.*, 68(4): 2386-9 (2000)).

Drug treatment

- Following one hour of acclimation to the observation cage, ferrets received
20 an intraperitoneal (i.p.) injection of cisplatin (5 mg/kg in 5 mL) followed in about 2 minutes by i.p. injection of a single dose of MCI-225 or ondansetron (Rudd and Naylor, *Eur. J. Pharmacol.*, 322: 79-82 (1997)). Dose-response effects of MCI-225 dosed at 1, 10 and 30 mg/kg i.p. in a 0.5 mL/kg solution or ondansetron dosed at 5 and 10 mg/kg i.p. in a 0.5 mL/kg solution were studied. Each animal received a
25 single-dose drug treatment. In addition, three animals received an initial dose (30 mg/kg i.p.) and a second MCI-225 injection (30 mg/kg i.p.) 180 minutes following the initial dose. Control animals were treated with cisplatin followed by vehicle alone (propanediol dosed in a 0.5 mL/kg solution). All groups were randomized.

Results

Vehicle Alone

Cisplatin induced an emetic response in 100 % of the animals receiving vehicle. The mean response was characterized by a total number of 42.8 ± 8.1 events (both retches and vomits), which occurred during the observation period. The mean latency of the first response was 133 ± 22 min post-cisplatin administration. The time-course of acute emetic events appearing in response to cisplatin is summarized in FIG. 1 (retches) and FIG. 2 (vomits).

Ondansetron

Ondansetron applied at the 5 mg/kg and 10 mg/kg dose-dependently reduced the number of emetic events induced by cisplatin. The effect of ondansetron was accompanied by an increase in the latency of the first emetic response following cisplatin treatment. The results are set forth in Table 1 (* $p < 0.05$), and presented graphically in FIG. 3 and FIG. 4.

TABLE 1

Nº. of Animal s (N)	Treatment	Retches (360 min)	Vomits (360 min)	Total Events	Latency (min)
N=10	Vehicle	42.8 ± 8.1	3.3 ± 0.8	46.1 ± 7.8	133 ± 22
N=7	Ondansetron (5 mg/kg)	11.2 ± 7.0	0.3 ± 0.2	11.5 ± 7.2	288 ± 4
N=7	Ondansetron (10 mg/kg)	2.4 ± 1.6	0.0 ± 0.0	$2.4 \pm 1.6^*$	313 ± 32

MCI-225

As set forth in Table 2, administration of MCI-225 at concentrations of 1, 10 or 30 mg/kg caused dose-dependent reduction in the retches and vomits induced by cisplatin (* $p < 0.05$). The emetic response was eliminated by administration of two doses of 30 mg/kg, applied b.i.d at a 180-min interval. The decrease in the number

of emetic events induced by MCI-225 was accompanied by an increase in the latency of the response. The results are set forth in Table 1 in FIG. 5 and FIG. 6.

TABLE 2

5	No. of Animals (N)	Treatment	Retches (360 min)	Vomits (360 min)	Total Events	Latency (min)
	N=10	Vehicle	42.8 ± 8.1	3.3 ± 0.8	46.1 ± 7.8	133 ± 22
	N=10	MCI-225 (1 mg/kg)	30.4 ± 9.1	2.5 ± 0.7	32.9 ± 9.8	186 ± 35
	N=10	MCI-225 (10 mg/kg)	22.9 ± 10.3	2.6 ± 1.0	$25.5 \pm$ 11.1	192 ± 57
10	N=11	MCI-225 (30 mg/kg)	3.3 ± 2.2	0.7 ± 0.5	$4.0 \pm 2.6^*$	287 ± 38
	N=3	MCI-225 (30 mg/kg b.i.d)	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	360 ± 0

Conclusion

The results set forth in Tables 1 and 2 show that MCI-225 is effective at reducing retching and vomiting in an accepted animal model of emesis, using a similar dose range as the positive control (ondansetron). Thus, MCI-225 can be used in the treatment of nausea, vomiting, retching or any combination thereof in a subject.


While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

-51-


CLAIMS

What is claimed is:

1. A method of treating nausea, vomiting, retching or any combination thereof
 5 in a subject in need thereof comprising administering to said subject a therapeutically effective amount of a compound of Formula I:



- 10 wherein, R_1 and R_2 independently represent hydrogen, halogen or a C_1 - C_6 alkyl group; or R_1 and R_2 together with the carbon atom to which they are attached form a cycloalkylene group having 5 to 6 carbon atoms;
 R_3 and R_4 independently represent hydrogen or a C_1 - C_6 alkyl group;
 R_5 is hydrogen, C_1 - C_6 alkyl,

or $-C(O)-NH-R_6$


wherein m is an integer from about 1 to about 3, X is halogen and R_6 is a C_1 - C_6 alkyl group; and

- Ar is a substituted or unsubstituted phenyl, 2-thienyl or 3-thienyl group; and
 20 n is 2 or 3; or a pharmaceutically acceptable salt thereof.

2. The method of Claim 1, wherein the the nausea, vomiting, retching or any combination thereof is caused by an anesthetic, radiation, a cancer chemotherapeutic agent, a toxic agent, an odor, a medicine, pregnancy or motion.
- 5 3. The method of Claim 2, wherein the medicine is selected from the group consisting of an analgesic, an antibiotic, an antifungal or a serotonin reuptake inhibitor.
4. The method of Claim 1, wherein the nausea, vomiting, retching or any combination thereof is caused by a condition which is associated with
10 vertigo.
5. The method of Claim 1, wherein the nausea, vomiting, retching or any combination thereof is caused by headache.
6. The method of Claim 1, wherein the nausea, vomiting, retching or any combination thereof is caused by a malady of the gastrointestinal (GI) tract.
- 15 7. The method of Claim 1, wherein the subject is a human.
8. The method of Claim 1, wherein for the compound of Formula I, R_1 is a C_1 - C_6 alkyl group and Ar is a substituted phenyl.
9. The method of Claim 8, wherein the substituted phenyl group is substituted
20 with a halogen.
10. The method of Claim 1, wherein for the compound of Formula I, n is 2, R_1 is a C_1 - C_6 alkyl group and Ar is a substituted phenyl.

-53-

11. The method of Claim 10, wherein the substituted phenyl group is substituted with a halogen and R_1 is a methyl group.
12. The method of Claim 1, wherein for the compound of Formula I, R_1 is a C_1 - C_6 alkyl group or a halogen and Ar is an unsubstituted phenyl.
- 5 13. The method of Claim 12, wherein R_2 is hydrogen or a C_1 - C_6 alkyl group.
14. The method of Claim 1, wherein for the compound of Formula I, n is 2, R_1 is a C_1 - C_6 alkyl group and Ar is an unsubstituted phenyl.
15. The method of Claim 14, wherein R_2 is hydrogen or a C_1 - C_6 alkyl group.
16. A method of treating nausea, vomiting, retching or any combination thereof
10 in a subject in need thereof comprising administering to said subject a therapeutically effective amount of a compound represented by Formula II:



II

or a pharmaceutically acceptable salt thereof.

17. The method of Claim 16, wherein the the nausea, vomiting, retching or any combination thereof is caused by an anesthetics, radiation, a cancer chemotherapeutic agent, a toxic agent, an odor, a medicine, pregnancy or motion.
- 5 18. The method of Claim 16, wherein the nausea, vomiting, retching or any combination thereof is caused by a condition which is associated with vertigo.
19. The method of Claim 16, wherein the nausea, vomiting, retching or any combination thereof is caused by headache.
- 10 20. The method of Claim 16, wherein the nausea, vomiting, retching or any combination thereof is caused by a malady of the gastrointestinal (GI) tract.
21. The method of Claim 16, wherein the subject is a human.
22. A method for treating nausea, vomiting, retching or any combinatin thereof in a subject in need thereof comprising administering to said subject:
- 15 a) a first amount of a 5-HT₃ receptor antagonist; and
b) a second amount of a noradrenaline reuptake inhibitor
wherein the first and second amounts together comprise a therapeutically effective amount.
- 20 23. The method of Claim 22, wherein the the nausea, vomiting, retching or any combination thereof is caused by an anesthetic, radiation, a cancer chemotherapeutic agent, a toxic agent, an odor, a medicine, pregnancy or motion.

24. The method of Claim 22, wherein the medicine is selected from the group consisting of an analgesic, an antibiotic, an antifungal or a serotonin reuptake inhibitor.
25. The method of Claim 22, wherein the nausea, vomiting, retching or any combination thereof is caused by a condition which is associated with vertigo.
26. The method of Claim 22, wherein the nausea, vomiting, retching or any combination thereof is caused by headache.
27. The method of Claim 22, wherein the nausea, vomiting, retching or any combination thereof is caused by a malady of the gastrointestinal (GI) tract.
28. The method of Claim 22, wherein the subject is a human.
29. The method of Claim 22, wherein the 5-HT₃ receptor antagonist is selected from the group consisting of indisetron, YM-114 ((R)-2,3-dihydro-1-[(4,5,6,7-tetrahydro-1H-benzimidazol-5-yl)-carbonyl]-1H-indole), granisetron, talipexole, azasetron, bemisetron, tropisetron, ramosetron, ondansetron, palonosetron, lerisetron, alosetron, N-3389, zacopride, cilansetron, E-3620 ([3(S)-endo]-4-amino-5-chloro-N-(8-methyl-8-azabicyclo[3.2.1]-oct-3-yl-2[(1-methyl-2-butynyl)oxy]benzamide), lintopride, KAE-393, itasetron, zatosectron, dolasetron, (±)-zacopride, (±)-renzapride, (-)-YM-060, DAU-6236, BIMU-8 and GK-128 [2-[2-methylimidazol-1-yl)methyl]-benzo[f]thiochromen-1-one monohydrochloride hemihydrate].

30. The method of Claim 29, wherein the 5-HT₃ receptor antagonist is selected from the group consisting of indisetron, granisetron, azasetron, bemisetron, tropisetron, ramosetron, ondansetron, palonosetron, lerisetron, alosetron, cilansetron, itasetron, zatoksetron, and dolasetron.
- 5 31. The method of Claim 22, wherein the noradrenaline reuptake inhibitor is selected from the group consisting of venlafaxine, duloxetine, bupropion, milnacipran, reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotline, viloxazine and
10 atomoxetine.
32. The method of Claim 31, wherein the noradrenaline reuptake inhibitor is selected from the group consisting of reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotline, viloxazine and atomoxetine.
- 15 33. A method for treating nausea, vomiting, retching or any combination thereof in a subject in need thereof comprising administering to said subject:
- a) a therapeutically effective amount of a 5-HT₃ receptor antagonist;
and
 - b) a therapeutically effective amount of a noradrenaline reuptake
20 inhibitor.
34. The method of Claim 33, wherein the the nausea, vomiting, retching or any combination thereof is caused by an anesthetic, radiation, a cancer chemotherapeutic agent, a toxic agent, an odor, a medicine, pregnancy or motion.
- 25 35. The method of Claim 33, wherein the medicine is selected from the group consisting of an analgesic, an antibiotic, an antifungal or a serotonin reuptake inhibitor.

-57-

36. The method of Claim 33, wherein the nausea, vomiting, retching or any combination thereof is caused by a condition which is associated with vertigo.
37. The method of Claim 33, wherein the nausea, vomiting, retching or any combination thereof is caused by headache.
38. The method of Claim 33, wherein the nausea, vomiting, retching or any combination thereof is caused by a malady of the gastrointestinal (GI) tract.
39. The method of Claim 33, wherein the subject is a human.
40. The method of Claim 33, wherein the 5-HT₃ receptor antagonist is selected from the group consisting of indisetron, YM-114 ((R)-2,3-dihydro-1-[(4,5,6,7-tetrahydro-1H-benzimidazol-5-yl)-carbonyl]-1H-indole), granisetron, talipexole, azasetron, bemisetron, tropisetron, ramosetron, ondansetron, palonosetron, lerisetron, alosetron, N-3389, zacopride, cilansetron, E-3620 ([3(S)-endo]-4-amino-5-chloro-N-(8-methyl-8-azabicyclo[3.2.1]-oct-3-yl-2[(1-methyl-2-butynyl)oxy]benzamide), linctopride, KAE-393, itasetron, zatosetron, dolasetron, (±)-zacopride, (±)-renzapride, (-)-YM-060, DAU-6236, BIMU-8 and GK-128 [2-[2-methylimidazol-1-yl)methyl]-benzo[f]thiochromen-1-one monohydrochloride hemihydrate].
41. The method of Claim 40, wherein the 5-HT₃ receptor antagonist is selected from the group consisting of indisetron, granisetron, azasetron, bemisetron, tropisetron, ramosetron, ondansetron, palonosetron, lerisetron, alosetron, cilansetron, itasetron, zatosetron, and dolasetron.

-58-

42. The method of Claim 33, wherein the noradrenaline reuptake inhibitor is selected from the group consisting of venlafaxine, duloxetine, bupropion, milnacipran, reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotline, viloxazine and atomoxetine.
43. The method of Claim 42, wherein the noradrenaline reuptake inhibitor is selected from the group consisting of reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotline, viloxazine and atomoxetine.
44. A method of treating nausea, vomiting, retching or any combination thereof in a subject in need thereof comprising administering to said subject a therapeutically effective amount of a noradrenaline reuptake inhibitor, wherein the noradrenaline reuptake inhibitor characterized by the substantial absence of anticholinergic effects.
45. The method of Claim 44, wherein the the nausea, vomiting, retching or any combination thereof is caused by an anesthetic, radiation, a cancer chemotherapeutic agent, a toxic agent, an odor, a medicine, pregnancy or motion.
46. The method of Claim 45, wherein the medicine is selected from the group consisting of an analgesic, an antibiotic, an antifungal or a serotonin reuptake inhibitor.
47. The method of Claim 44, wherein the nausea, vomiting, retching or any combination thereof is caused by a condition which is associated with vertigo.

-59-

48. The method of Claim 44, wherein the nausea, vomiting, retching or any combination thereof is caused by headache.
49. The method of Claim 44, wherein the nausea, vomiting, retching or any combination thereof is caused by a malady of the gastrointestinal (GI) tract.
50. The method of Claim 44, wherein the subject is a human.
51. The method of Claim 44, wherein the noradrenaline reuptake inhibitor is selected from the group consisting of venlafaxine, duloxetine, bupropion, milnacipran, reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotline, viloxazine and atomoxetine.
52. The method of Claim 51, wherein the noradrenaline reuptake inhibitor is selected from the group consisting of reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotline, viloxazine and atomoxetine.
53. A pharmaceutical composition comprising:
- a first amount of a 5-HT₃ receptor antagonist; and
 - a second amount of a noradrenaline reuptake inhibitor.
54. The pharmaceutical composition of Claim 53, further comprising a pharmaceutically acceptable carrier.
55. The pharmaceutical composition of Claim 53, wherein the 5-HT₃ receptor antagonist is selected from the group consisting of indisetron, YM-114 ((R)-2,3-dihydro-1-[(4,5,6,7-tetrahydro-1H-benzimidazol-5-yl)-carbonyl]-1H-indole), granisetron, talipexole, azasetron, bemesetron, tropisetron, ramosetron, ondansetron, palonosetron, lerisetron, alosetron, N-3389,

-60-

zacopride, cilansetron, E-3620 ([3(S)-endo]-4-amino-5-chloro-N-(8-methyl-8-azabicyclo[3.2.1]-oct-3-yl-2-[(1-methyl-2-butynyl)oxy]benzamide), linctopride, KAE-393, itasetron, zatosetron, dolasetron, (\pm)-zacopride, (\pm)-renzapride, (-)-YM-060, DAU-6236, BIMU-8 and GK-128 [2-[2-methylimidazol-1-yl)methyl]-benzo[f]thiochromen-1-one monohydrochloride hemihydrate].

56. The pharmaceutical composition of Claim 55, wherein the 5-HT₃ receptor antagonist is selected from the group consisting of indisetron, granisetron, azasetron, bemisetron, tropisetron, ramosetron, ondansetron, palonosetron, lerisetron, alosetron, cilansetron, itasetron, zatosetron, and dolasetron.
57. The pharmaceutical composition of Claim 53, wherein the noradrenaline reuptake inhibitor is selected from the group consisting of venlafaxine, duloxetine, bupropion, milnacipran, reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotiline, viloxazine and atomoxetine.
58. The pharmaceutical composition of Claim 53, wherein the noradrenaline reuptake inhibitor is selected from the group consisting of reboxetine, lefepramine, desipramine, nortriptyline, tomoxetine, maprotiline, oxaprotiline, levoprotiline, viloxazine and atomoxetine.
59. A method for processing a claim under a health insurance policy submitted by a claimant seeking reimbursement for costs associated with treatment of nausea, vomiting, retching or any combination thereof, wherein said treatment comprises coadministering to a subject a first amount of a 5-HT₃ receptor antagonist and a second amount of a noradrenaline reuptake inhibitor, wherein the first and second amounts together comprise a therapeutically effective amount comprising:
- a) reviewing said claim;

-61-

- b) determining whether said treatment is reimbursable under said insurance policy; and
- c) processing said claim to provide partial or complete reimbursement of said costs.

5 60. The method of Claim 59, wherein the nausea, vomiting, retching or any combination thereof is caused by an anesthetic, radiation, a cancer chemotherapeutic agent, a toxic agent, an odor, a medicine, pregnancy or motion.

10 61. The method of Claim 60, wherein the medicine is selected from the group consisting of an analgesic, an antibiotic, an antifungal or a serotonin reuptake inhibitor.

62. The method of Claim 59, wherein the nausea, vomiting, retching or any combination thereof is caused by a condition which is associated with vertigo.

15 63. The method of Claim 59, wherein the nausea, vomiting, retching or any combination thereof is caused by headache.

64. A method of Claim 59, wherein the nausea, vomiting, retching or any combination thereof is caused by a malady of the gastrointestinal (GI) tract.

20 65. A method for processing a claim under a health insurance policy submitted by a claimant seeking reimbursement for costs associated with treatment of nausea, vomiting, retching or any combination thereof, wherein said treatment comprises coadministering to a subject a therapeutically effective amount of a 5-HT₃ receptor antagonist and a therapeutically effective amount of a noradrenaline reuptake inhibitor comprising:

25 a) reviewing said claim;

-62-

- b) determining whether said treatment is reimbursable under said insurance policy; and
- c) processing said claim to provide partial or complete reimbursement of said costs.

- 5 66. The method of Claim 65, wherein the nausea, vomiting, retching or any combination thereof is caused by an anesthetic, radiation, a cancer chemotherapeutic agent, a toxic agent, an odor, a medicine, pregnancy or motion.
- 10 67. The method of Claim 66, wherein the medicine is selected from the group consisting of an analgesic, an antibiotic, an antifungal or a serotonin reuptake inhibitor.
68. The method of Claim 65, wherein the nausea, vomiting, retching or any combination thereof is caused by a condition which is associated with vertigo.
- 15 69. The method of Claim 65, wherein the nausea, vomiting, retching or any combination thereof is caused by headache.
70. The method of Claim 65, wherein the nausea, vomiting, retching or any combination thereof is caused by a malady of the gastrointestinal (GI) tract.

1/3




FIG. 1




FIG. 2

2/3




FIG. 3




FIG. 4

3/3




FIG. 5




FIG. 6